



ASI SNAKE BOOKLET

COMMON SNAKES OF BOTSWANA

Copyright: Johan Marais - African Snakebite Institute
First Edition published 2025

814 Wapadrand Rd,
Wapadrand Security Village,
Pretoria, 0081

Cell: +27 (0)82 494 2039
E-mail johan@asiorg.co.za
www.africansnakebiteinstitute.com

All rights reserved. No part of this publication may be reproduced or used commercially without the prior written permission of the copyright owner.

SNAKE COURSES

The ASI offers both public and corporate snake courses.

Public courses take place at various venues throughout South Africa.

The corporate training is done on-site, both in South Africa, as well as in 19 African countries.

We pride ourselves on teaching only the safest snake handling methods, to ensure the safety of both the handler and the snake.

We offer a range of different course options, including snake awareness, first aid for snakebite, venomous snake handling, venomous snake handling bootcamps, as well as advanced first aid for snakebite, advanced venomous snake handling and a reptile photography course.

For course info,
click [here](#) or email us on
courses@asiorg.co.za
for details.



www.AFRICANSNAKEBITEINSTITUTE.com



INTRODUCTION

The African Snakebite Institute is the leading training provider of Snake Awareness, First Aid for Snakebite and Venomous Snake Handling courses in Africa as well as the largest distributor of quality snake handling equipment, snake PPE and snakebite first aid kits. Our aim is to spread snake awareness as far and wide as possible, which is why we've launched our new free E-Book series.

Our CEO, Johan Marais is a well-known herpetologist with over 40 years of experience with snakes. He has worked at various reptile facilities and has collaborated on a variety of scientific studies with some of the world's leading herpetologists. He is also one of Africa's most prominent authors of snake books and has written the best-sellers *A Complete Guide to Snakes of Southern Africa*, *Snakes and Snakebite in Southern Africa* and *First Aid for Snakebite*, to mention just a few.



COMMON SNAKES OF BOTSWANA

Botswana has 62 different snake species. Of these, eight are highly venomous, ten can give a bad bite and the rest are mildly venomous or harmless. We've listed the common species in this booklet, showing venomosity, key ID features, full distribution, and snakebite info for venomous species.

Snakebite is problematic in Southern Africa with most bites reported from the hot and humid regions. Most bites are caused by snakes with cytotoxic venom. These bites can result in extensive tissue damage. Snakes responsible for these type of bites include the Puff Adder, Night Adders and the Spitting Cobras. Although not as frequent, neurotoxic bites, inflicted by non-spitting cobras and Mambas can pose serious medical emergencies as patients may stop breathing.

Most bites occur in the early evening in the warm, wet summer months from November to April/May. People most at risk are those that spend a lot of time outdoors. Agricultural workers in rural areas are especially at risk. Most snakebites are inflicted on the leg below the knee. This is as a result of people accidentally stepping onto or close to snakes. Given that a patient receives prompt and proper treatment in a hospital the chances of surviving a snakebite are very high. In a study performed at a clinic in Kwa Zulu Natal in South Africa not one patient died during the five year study thanks to the treatment they received. The clinic treated 879 patients for snakebites in those five years!

SNAKE HANDLING EQUIPMENT

The African Snakebite Institute is the leading provider of quality snake handling equipment in Africa. View our range online [here](http://www.AFRICANSNAKEBITEINSTITUTE.com).



www.AFRICANSNAKEBITEINSTITUTE.com

COMMON SNAKES OF BOTSWANA



Botswana has 62 species of snakes. Eight species are considered potentially deadly. Another ten species can give a bad bite. Twenty species are mildly venomous and not dangerous to humans. Twenty-four species are harmless.

HIGHLY VENOMOUS

Has caused human fatalities

VENOMOUS

Painful bite, but does not require antivenom

MILDLY VENOMOUS

Not thought to be harmful

HARMLESS

Not dangerous to humans



Black Mamba
(*Dendroaspis polylepis*)

HIGHLY VENOMOUS



Boomslang - male
(*Dispholidus typus*)

HIGHLY VENOMOUS



Southern Vine Snake
(*Thelotornis capensis*)

HIGHLY VENOMOUS



Spouted Cobra
(*Naja annulifera*)

HIGHLY VENOMOUS



Puff Adder
(*Bitis arietans*)

HIGHLY VENOMOUS



Mozambique Spitting Cobra
(*Naja mossambica*)

HIGHLY VENOMOUS



Cape Cobra
(*Naja nivea*)

HIGHLY VENOMOUS



Anchieta's Cobra
(*Naja anchietae*)

HIGHLY VENOMOUS



Speckled Shield-nose Snake
(*Aspidelaps x. scutatus*)

VENOMOUS



Bibron's Stiletto Snake
(*Atractaspis bibronii*)

VENOMOUS



Southern African Python
(*Python natalensis*)

CAN INFLICT A NASTY BITE



Mole Snake
(*Pseudaspis cana*)

CAN INFLICT A NASTY BITE



Olive Grass Snake
(*Psammophis mossambicus*)

MILDLY VENOMOUS



Western Yellow-bellied Sand Snake
(*Psammophis subtaeniatus*)

MILDLY VENOMOUS



Eastern Tiger Snake
(*Teloscopus semiannulatus*)

MILDLY VENOMOUS



Striped Skaapsteker
(*Psammophylax triaenatus*)

MILDLY VENOMOUS



Herald Snake
(*Crotaphopeltis hotamboeia*)

MILDLY VENOMOUS



Spotted Bush Snake
(*Philothamnus semivariegatus*) - photo Luke Kemp

HARMLESS



Brown House Snake
(*Boaedon capensis*)

HARMLESS



Common Egg-eater
(*Dasypeltis scabra*)

HARMLESS

JOHAN MARAIS is the author of various books on reptiles including the best-seller *A Complete Guide to Snakes of Southern Africa*. He is a popular public speaker and CEO of the African Snakebite Institute. The African Snakebite Institute (ASI) offers a variety of courses including Snake Awareness, Scorpion Awareness and Venomous Snake Handling. ASI is accredited by the International Society of Zoological Sciences (ISZS) and Travel Doctor-approved service provider. The courses are also accredited by the Health Professions Council of South Africa (HPCSA). ASI is the largest provider of quality snake handling and safety equipment in Africa.

Johan Marais | African Snakebite Institute

+27 82 494 2039 | johan@asiorg.co.za

www.AFRICANSNAKEBITEINSTITUTE.com

Get the **FREE ASI SNAKES APP**

Use our app to learn about snakes, find trained snake removers in your area, browse our free resources, and more.



Download the high resolution poster [here](#).

www.AFRICANSNAKEBITEINSTITUTE.com



HIGHLY VENOMOUS



Black Mamba

(Dendroaspis polylepis)

Maximum Length:
3.8 m



Other names:

Swartmamba

This is by far the largest venomous snake in Africa, with a maximum length of 3.8 m (historically specimens over 4 m were reported). Despite its reputation it is a shy, elusive snake that is quick to escape but will not hesitate to strike repeatedly if cornered. It is active during the day, often basking near a hole in an anthill or large rock crevice, into which it quickly disappears if disturbed. When threatened it will gape exposing the black inner lining of the mouth and it may form a narrow hood. The venom of this snake is potently neurotoxic and may cause difficulty with breathing within half an hour. Antivenom is effective but often required in large quantities (10 – 15 vials).



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Black Mamba (*Dendroaspis polylepis*)

HIGHLY VENOMOUS

Average length: 2.2 - 2.6 m. Maximum length: 4.5 m

Can be confused with: Boomslang, Olive Grass Snakes, Cobras



Coffin-shaped head

Can lift up to a third of its body off the ground

Colour varies from light olive to light grey, darkening with age to olive, greyish brown, or gunmetal grey. Sometimes has mottling

Elongated coffin-shaped head

Inside of the mouth is usually black

Forms a narrow hood when threatened

May have barring toward the tail

Light grey belly, sometimes with mottling


www.AFRICANSNAKEBITEINSTITUTE.com



SNAKEBITE PROFILE

BLACK MAMBA

Dendroaspis polylepis



Risk of Bites

The Black Mamba is a nervous snake and is quick to escape. If accidentally cornered in a confined space or surprised in the veld, it may strike out and bite more than once in quick succession. Contrary to popular stories, it does not chase people and bites are quite rare.



Most Bites

Most victims are bitten in the veld when they accidentally get very close to a Black Mamba basking. Or when accidentally cornering a mamba in a confined space like a pump house on a farm. Never approach a Black Mamba closely and do not try to kill or catch one.



Symptoms & Venom

A potent and fast-acting neurotoxic venom that results in pins and needles in the lips, hands and feet, as well as nausea, excessive sweating, drooping eyelids, difficulty in swallowing and eventually drowsiness and muscle pain followed by paralysis and respiratory failure. In severe cases victims become unresponsive within less than one hour.

See pages 43-45

– *Snakes and Snakebite in Southern Africa* (2024).




First Aid

Immobilise and reassure the patient, who must lie down in a rescue position. Immediately arrange transport to a hospital with emergency facilities. If far from a hospital, apply a pressure bandage (Page 19 of *Snakes and Snakebite in Southern Africa*). Resort to artificial respiration if the victim stops breathing.



Medical Treatment

In cases of severe envenomation where patients are not breathing, they are intubated and ventilated and treated with 12-15 vials of SAVP Polyvalent antivenom. Additional vials of antivenom may be required at a later stage.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




HIGHLY VENOMOUS



Snouted Cobra (*Naja annulifera*)

Maximum Length:
2.5m



Other names:

Wipneuskobra, Bosveldkapel

The Snouted Cobra is largely a moist and dry savannah inhabitant but is also found in lowveld and bushveld areas where it actively hunts its prey during the day or at night. It feeds on rodents, birds, their eggs and especially toads, but is fond of eating snakes, including the Puff Adder. It is not an aggressive snake but if cornered will stand its ground and form an impressive hood. It may exceed 2.5 m in length. It is usually different shades of brown, but some individuals have broad light brown and purplish-black bands on the back. The venom of this snake is both neurotoxic and cytotoxic and may cause some tissue damage. Polyvalent antivenom is effective, but luckily bites are rare.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com


www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID


Snouted Cobra (*Naja annulifera*)

HIGHLY VENOMOUS



Average length: 1.2 - 1.6 m. Maximum length: 2.5 m

Can be confused with: Mozambique Spitting Cobras, Cape cobras



www.AFRICANSNAKEBITEINSTITUTE.com



Snakebite Profile

Snouted Cobra

Naja annulifera



Risk of Bites

Bites from the Snouted Cobra are not common. This snake frequently visits residential gardens in search of rodents and toads. It also raids poultry runs. Dogs are at risk and often get bitten when attempting to kill the snake.




Most Bites

Bites are infrequent and usually occur when accidentally confronted or stepped on at night. Most bites are on the lower legs, or to those who try and handle the snake.




Symptoms & Venom

The venom is both neurotoxic and cytotoxic. It is one of our largest cobras reaching 2,5 m in length and can deliver a lot of venom in a single bite. Bites may result in burning pain, nausea and swelling followed by tissue damage. It can also affect breathing.



See pages 52-53

– *Snakes and Snakebite in Southern Africa* (2024).



First Aid

Immobilise and reassure the patient, who must lie down in a rescue position. Immediately arrange transport to a hospital with emergency facilities. If far from a hospital, apply a pressure bandage (Page 19 of *Snakes and Snakebite in Southern Africa*). Resort to artificial respiration if the victim stops breathing.



Medical Treatment

In cases of severe envenomation, especially where patients are not breathing, they should be intubated, ventilated, and treated with 12-15 vials of SAVP Polyvalent antivenom. Additional vials of antivenom may be required at a later stage.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



HIGHLY VENOMOUS



Anchieta's Cobra (*Naja anchietae*)

Maximum Length:
2.5 m



Other names:

Anchieta se Kobra

Anchieta's Cobra is very similar to the Snouted Cobra in both appearance and behaviour. It occurs in western Zimbabwe, northern Botswana and northern Namibia (and elsewhere further north). Individuals are orange-brown to purple-brown and banded varieties are also found. This snake has 17 midbody scale rows (the Snouted Cobra has 19) and the scales appear much larger than in the Snouted Cobra. Maximum size is around 2.5 m. It seldom features in snakebite incidents but will spread an impressive hood if threatened. The venom of this snake is predominantly neurotoxic but may also cause blistering and tissue damage. Because of its size a bite from Anchieta's Cobra could result in alarming symptoms. Antivenom is effective against its bite.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com





ASI EASY ID

Anchieta's Cobra (*Naja anchietae*)

HIGHLY VENOMOUS

Average length: 1.2 - 1.5 m. Maximum length: 2.4 m

Often confused with: Snouted Cobras



www.AFRICANSNAKEBITEINSTITUTE.com


www.AFRICANSNAKEBITEINSTITUTE.com



SNAKEBITE PROFILE

ANCHIETA'S COBRA

Naja anchietae



Risk Of Bites

Bites are not common, although this snake may be frequently found around human dwellings. It is also known to raid poultry runs. Dogs are at risk and often get bitten when attempting to kill the snake.



Most Bites

Bites are infrequent and usually occur when accidentally confronted or stepped on at night. Most bites are on the lower legs, or to those who try and handle the snake.



Symptoms & Venom

The venom is both neurotoxic and cytotoxic. It is one of our largest cobras reaching 2.5m in length and it can deliver large volumes of venom in a single bite. Bites may result in burning pain, nausea and swelling followed by tissue damage. It can also result in progressive weakness and affect breathing.

See pages 52-53

– *Snakes and Snakebite in Southern Africa* (2024).




First Aid

Immobilise and reassure the patient, who must lie down in a rescue position. Immediately arrange transport to a hospital with emergency facilities. If far from a hospital, apply a pressure bandage (Page 19 of *Snakes and Snakebite in Southern Africa*). Resort to artificial respiration if the victim stops breathing.



Medical Treatment

In cases of severe envenomation, especially where patients are not breathing, they should be intubated, ventilated and treated with 12-15 vials of SAVP Polyvalent antivenom. Additional vials of antivenom may be required at a later stage.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




HIGHLY VENOMOUS



Cape Cobra

(*Naja nivea*)

Maximum Length:
2.3 m



Other names:

Geelslang, Kaapse Kobra, Koper kapel

This snake varies in colour from near black to dark or light brown, beige, yellow or speckled while juveniles have a dark band on the throat. The Cape Cobra is easily confused with the Mole Snake and the Black Spitting Cobra. It is by far our most dangerous cobra and with the Black Mamba it accounts for the majority of fatal snake bites in South Africa. The Cape Cobra may stand its ground if threatened and is quick to form a hood and bite. It occurs largely in the Cape provinces but extends into the Free State, Northwest, southern Botswana and Namibia. The venom is potently neurotoxic causing progressive weakness and may affect breathing rapidly. Victims need urgent hospitalisation and in a severe bite antivenom is essential.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID


Cape Cobra (*Naja nivea*)

HIGHLY VENOMOUS



Average length: 1.2 - 1.6 m. Maximum length: 2.3 m

Can be confused with: Mole Snakes, Black Spitting Cobras, Snouted Cobras




Often has a dark mark under the eye

Chunky cheeks for venom storage

Varies in colour from light yellow to brown, gold and black.
Often a speckled mix of colours

Spreads an impressive hood

Juveniles have a dark band on throat



Ventral scales often visible on sides

Belly usually same colour as the rest of snake

Usually have dark tipped tails

www.AFRICANSNAKEBITEINSTITUTE.com

Snakebite Profile

CAPE COBRA

Naja nivea



Risk of Bites

The Cape Cobra is commonly encountered in suburbia, especially in areas where rodents are abundant. If cornered, it is quick to rear up and form a hood and if approached closely will bite readily.



Most Bites

Most victims are bitten on the lower leg or on a hand, often when attempting to capture or kill the snake.




Symptoms & Venom

A potent and fast-acting neurotoxic venom that is quite similar to Black Mamba venom in its effects. It results in pins and needles in the lips, nausea, excessive sweating, drooping eyelids, difficulty in swallowing and eventually drowsiness and muscle pain followed by paralysis and respiratory failure. Some victims also experience some swelling. In severe cases victims become unresponsive within less than one hour.

See pages 49-51

– *Snakes and Snakebite in Southern Africa* (2024).




First Aid

Immobilise and reassure the patient, who must lie down in a rescue position. Immediately arrange transport to a hospital with emergency facilities. If far from a hospital, apply a pressure bandage (Page 19 of *Snakes and Snakebite in Southern Africa*). Resort to artificial respiration if the victim stops breathing.



Medical Treatment

In cases of severe envenomation patients are intubated and ventilated if they are not breathing. They are treated with 12-15 vials of SAVP Polyvalent antivenom. Additional vials of antivenom may be required at a later stage. Even after receiving antivenom, patients may remain unresponsive for a day or two.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




HIGHLY VENOMOUS



Mozambique Spitting Cobra (*Naja mossambica*)

Maximum Length:
1.8 m



Other names:

Mosambiekse Spoegkobra

The Mozambique Spitting Cobra occurs on the eastern side of South Africa, most of Mozambique, Zimbabwe, north-eastern Botswana, northern Namibia and further north in Africa. Its diet consists largely of frogs, small mammals, birds and snakes, including the Puff Adder. It is active on overcast days, but more active at night, often ending up in houses where people are bitten while asleep. This snake accounts for the majority of serious snakebite cases in southern Africa, but fatalities are rare. It also spits its venom and does not always spread a hood when doing so. The venom is potentially cytotoxic causing pain, swelling, blisters and in many cases severe tissue damage. Antivenom is effective if administered early and in large dosages.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Mozambique Spitting Cobra (*Naja mossambica*)

HIGHLY VENOMOUS

Average length: 80 cm - 1.2 m. Maximum length: 1.83 m

Can be confused with: Rinkhals, Snouted Cobras



Black markings on the edges of the scales of the upper lip

Can spit effectively up to 3 m

Salmon coloured throat

Black bars and blotches on the throat

Spreads a hood, but can spit from any position

Slate grey to olive brown body

Belly may be off-white, to salmon in colour

Black edged scales

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



SNAKEBITE PROFILE

MOZAMBIQUE SPITTING COBRA

Naja mossambica



Risk of Bites

Bites are common, as this snake is widespread throughout much of southern Africa. Along with the Puff Adder, these two snakes account for most serious snakebites in southern Africa. Dogs are at risk and often get bitten when attempting to kill the snake.




Most Bites

Most victims are bitten at night when accidentally stepping on the snake, or while in bed. The snake enters homes and may bite sleeping victims, often multiple times. Bites in bed appear to be somewhat of a feeding response with the snake smelling a mammal and biting. Should you live within the range of this snake it is advisable to ensure that exit doors seal well without gaps or to fix mosquito-proof doors to all exits. People are also encouraged to sleep under mosquito nets.



Symptoms & Venom


The venom is predominantly and potently cytotoxic. Symptoms include severe pain, progressive swelling, various lesions and, in some cases, severe tissue damage.



www.AFRICANSNAKEBITEINSTITUTE.com

See pages 64-65

– *Snakes and Snakebite
in Southern Africa*
(2024).



First Aid

Immobilise and reassure the patient. Elevate the affected limb and remove tight clothing and jewelry. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages. For venom in the eyes, wash the eyes out immediately using water or any other bland liquid and get the victim to a medical doctor for further examination and treatment.



Medical Treatment

In cases of severe envenomation where there are clear signs of progressive swelling, extending more than 10-15cm per hour above the bite site, doctors must treat such bites with 12 ampoules (or more) of polyvalent antivenom. It appears that SAVP polyvalent antivenom is not very effective against the venom of this snake, therefore it should be administered within an hour or two after a bite. Anti-inflammatories (NSAID's) and antibiotics are not indicated in the short term. Fasciotomies are rarely justified and should generally be avoided. In cases where patients suffer tissue damage, surgery may be required but rarely in the first few days.

Fatalities are rare.






HIGHLY VENOMOUS



Boomslang (*Dispholidus typus*)

Maximum Length:
2.1 m



Other names:

Boomslang

The Boomslang is largely tree-living but may descend to the ground to bask. In trees it poses no threat to humans as it is extremely reluctant to bite, and bites are rare. Juveniles are grey with a massive emerald, green eye but change to the adult colours around a length of 1 m. Most males are bright green, sometimes with black between the scales, but in the Cape Provinces they are usually black above with green, yellow or orange sides. Females are usually brown to grey in colour. Boomslang venom is haemotoxic and compromises the blood clotting mechanism, causing uncontrolled bleeding if not treated. The South African Vaccine Producers manufacture a monovalent antivenom for bites from Boomslang that is very effective.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Boomslang (*Dispholidus typus*)


HIGHLY VENOMOUS

Average length: 1.2 m. Maximum length: 2.1 m

Often confused with: Black Mambas, Green Mambas, green snakes and sand snakes




Females are normally olive grey to brown or brick red in colour



Small head with large eye


Scales are keeled giving snake a rough texture



Most males are green in colour



Males in the Cape are usually black above with yellow, green or orange sides



Large eye

Short head



Juveniles have big emerald green eyes

www.AFRICANSNAKEBITEINSTITUTE.com



SNAKEBITE PROFILE

BOOMSLANG

Dispholidus typus



Risk of Bites

Bites are not common in South Africa. This snake lives in trees and shrubs and interactions with humans are infrequent. It is a placid snake that only bites when provoked or handled. If spotted, it is quick to flee higher up into trees.



Most Bites

Bites occur when a basking snake is accidentally stepped on or grabbed, or when it is handled.




Symptoms & Venom

Drop for drop the Boomslang has the most potent venom of any snake in Africa. It is slow-acting, but potently haemotoxic and symptoms include bleeding from the bite site, the nose, mucous membranes and internal bleeding.

See pages 79-81

– *Snakes and Snakebite in Southern Africa* (2024).




First Aid

Immobilise and reassure the patient. Elevate the affected limb. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages.



Medical Treatment

In cases of severe envenomation patients are treated with monovalent Boomslang antivenom. The initial dosage is two vials but, in some cases, a third vial may be required. This antivenom is produced by the South African Vaccine Producers. It is a Schedule 4 drug, therefore a script is required should one wish to purchase it. It comes in a 10 ml vial and must be refrigerated. It has a three year shelf life.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




HIGHLY VENOMOUS



Southern Vine Snake (*Thelotornis capensis*)

Maximum Length:
1.47 m



Other names:

Twig Snake; Bird Snake; Takslang; Voëlslang

This perfectly-camouflaged tree-living snake is seldom seen because of its excellent camouflage and habit of remaining very still in low shrubs, observing the ground below for passing lizards and snakes. Birds often mob this snake, and it will inflate its neck with its bright orange tongue flickering – this leads to the incorrect assumption that they lure birds with their tongue. It is extremely placid but, if provoked, will inflate its neck and strike viciously. Most bites are to snake handlers. Like the Boomslang, the venom of this snake is haemotoxic affecting the blood clotting mechanism and causing uncontrolled bleeding. There is no antivenom for the venom of this snake and although a few fatalities have been reported, none were in South Africa.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com





ASI EASY ID

Southern Vine Snake (*Thelotornis c. capensis*)

HIGHLY VENOMOUS

Average length: 1.2 m. Maximum length: 1.47 m

Often confused with: Juvenile Boomslang, Eastern Bark Snakes



www.AFRICANSNAKEBITEINSTITUTE.com



SNAKEBITE PROFILE

SOUTHERN VINE SNAKE

Thelotornis capensis



Risk Of Bites

This snake lives in trees and shrubs and interactions with humans are not common. It is a placid snake that only bites when provoked or handled. Bites are very rare in South Africa.



Most Bites

There are very few documented bites from this snake in southern Africa, except for incidents where snake keepers are bitten by captive snakes, or by other people handling this snake.




Symptoms & Venom

The venom is potently haemotoxic, like that of the Boomslang. Symptoms include bleeding from the nose, mucous membranes and bite site. This may eventually lead to internal bleeding.

See pages 82-84

– *Snakes and Snakebite in Southern Africa* (2024).




First Aid

Immobilise and reassure the patient. Elevate the affected limb. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages.



Medical Treatment

Polyvalent and Monovalent antivenom do not offer cross coverage for Vine Snake bites and should not be administered. Bites are treated symptomatically with blood transfers and platelets, but this may not be effective in severe bites. To date we do not have any documented fatalities in South Africa, however a few fatal bites have been recorded further north in Africa.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



HIGHLY VENOMOUS



Puff Adder

(*Bitis arietans*)

Maximum Length:
1.4 m



Other names:

Pofadder, Imbululu

The Puff Adder is widespread over much of southern Africa and further north into Africa. It is a heavy-bodied snake that relies on its excellent camouflage and is reluctant to move. Much of its life is spent in camouflage mode and recent research has shown that when hiding, the Puff Adder will not hiss or strike when approached, as this would give away its presence. It is an ambush hunter that will coil up into a striking position and await its prey. This can last several days or even weeks. Toads are lured closer with flicking of the tongue. Puff Adder venom is potently cytotoxic, causing severe pain, swelling, blistering and in many cases severe tissue damage. Polyvalent antivenom is effective.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Puff Adder (*Bitis arietans*)

HIGHLY VENOMOUS

Average length: 90 cm. Maximum length: 1.4 m

Often confused with: night adders and Berg Adders



www.AFRICANSNAKEBITEINSTITUTE.com



SNAKEBITE PROFILE

PUFF ADDER

Bitis arietans



Risk Of Bites

Bites are common as this snake is widespread throughout most of southern Africa. Along with the Mozambique Spitting Cobra, these two snakes account for the majority of serious snakebites in southern Africa. Dogs are at risk and often get bitten when attempting to kill the snake.



Most Bites


Most victims are bitten at night when accidentally stepping on the snake, but bites are also inflicted during the day when the snake is basking. Most bites are on the feet or lower legs.



Symptoms & Venom

The venom is potently cytotoxic as well as haemotoxic. Symptoms include severe pain, progressive swelling, bleeding from the fang punctures, blistering and tissue damage. The lymph nodes may also be affected. Dogs may perish after a bite from this snake, unless treated with polyvalent antivenom, if required.

See pages 26-28
– *Snakes and Snakebite
in Southern Africa*
(2024).



First Aid

Immobilise and reassure the patient. Elevate the affected limb and remove tight clothing and jewelry. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages.



Medical Treatment

In cases of severe envenomation where there are clear signs of progressive swelling, which extends more than 10-15cm per hour above the bite site, doctors will treat such bites with 6 ampoules (or more) of polyvalent antivenom. Anti-inflammatories (NSAID's) and antibiotics are not indicated in the short term. Fasciotomies are rarely justified and should generally be avoided. In cases where patients suffer tissue damage, surgery may be required but rarely in the first few days.

Fatalities are rare.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




VENOMOUS



Speckled Shield-nose Snake (*Aspidelaps s. scutatus*)

Maximum Length:
65 cm



Other names:

Skildneusslang, Skildneuskobra

A smallish nocturnal snake that averages 40 – 45 cm in length and with a maximum length of 65 cm. It spends most of its time foraging in loose sand in search of small mammals, frogs and especially legless lizards, using its large rostral scale as a bulldozer.

If threatened it will lift its head off the ground, flatten the neck region into somewhat of a hood and strike repeatedly while hissing.

The venom of the Speckled Shield-nose Snake is mildly cytotoxic and neurotoxic, causing local swelling and necrosis. Respiratory failure may occur but is rare. There is no antivenom for the bite of this snake and patients must be treated symptomatically.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Speckled Shield-nose Snake (*Aspidelaps scutatus scutatus*)


VENOMOUS

Average length: 45 cm. Maximum length: 75 cm

Often confused with: other shield-nose snakes



Distribution
is important



Large rostral
scale

Black head

White patch
on throat


Black neck

Keeled scales on the back
half of body

Faint dark blotches
down spine

Reddish-brown
to grey in colour

Yellowish-white
to pink belly



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com

Snakebite Profile

Speckled Shield-nose Snake

Aspidelaps scutatus scutatus

Risk of Bites

Bites to humans are not common. This snake puts up a big show when cornered and is quick to hiss and strike repeatedly.


Most Bites

Bites occur mostly at night, when the snake is trodden on, or accidentally cornered. There is one recorded death of a child.

Symptoms & Venom

The venom of this snake is both cytotoxic and neurotoxic, resulting in local swelling and possibly necrosis. Paralyzed facial muscles, drooping eyelids (ptosis) and blurred vision have been recorded. Victims may experience breathing issues in severe bites.

See pages 74-75
– *Snakes and Snakebite
in Southern Africa*
(2024).




First Aid

Immobilise and reassure the patient and elevate the affected limb. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages.

Medical Treatment

There is no antivenom for bites from this snake and doctors will treat the bite symptomatically.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



VENOMOUS



Bibron's Stiletto Snake

(Atractaspis bibronii)

**Maximum Length:
70 cm**



Other names:

**Bibron se Sypikslang; Moladder; Mole
Viper; Burrowing Asp**

After the Mozambique Spitting Cobra and Puff Adder, the Bibron's Stiletto Snake accounts for most serious snakebites in South Africa. It spends most of its life underground but often emerges on hot humid nights and may be stepped on. With large hinged fangs it cannot be handled safely and also cannot be held behind the head – it merely twists its head sideways and will get a fang to penetrate a fingertip. It is usually blackish brown in colour and is easily confused for a variety of harmless snakes. This snake averages 30 cm but can exceed 70 cm in length. Its venom is potently cytotoxic causing severe pain, swelling, blistering and necrosis. Antivenom is not available for this snake.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Bibron's Stiletto Snake (*Atractaspis bibronii*)

VENOMOUS

Average length: 40 - 60 cm. Maximum length: 98 cm

Often confused with: wolf snakes, blind snakes, purple-glossed snakes



Body is brown to blackish in colour

Short stubby tail with spike on the end

Tries to bury head in sand, forming an arch in its neck

Has single subcaudal scales

Belly may be white or same colour as body

Small pig-like eyes

Large hinged fangs

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



Snakebite Profile

BIBRON'S STILETTO SNAKE

Atractaspis bibronii



Risk of Bites

Bibron's Stiletto Snake is largely active at night, especially on warm summer nights after a bout of rain. It may also fall into swimming pools and may be brought indoors by cats.



Most Bites

Most victims are bitten picking up Stiletto Snakes, mistaking them for a harmless snake. Otherwise, when stood on while walking barefoot at night.




Symptoms & Venom

The venom of this snake is potently cytotoxic, causing severe pain, local swelling, blistering, and potentially local tissue damage.

See pages 85-87

– *Snakes and Snakebite in Southern Africa* (2024).




First Aid

There is virtually nothing that can be done. Elevate the affected limb and get to a hospital that has an emergency department.



Medical Treatment

As there is no antivenom for a bite from this snake, doctors need to treat for pain and rehydrate the victim. Anti-inflammatories, antibiotics and antihistamines are not effective and should not be administered. It is equally as important not to resort to any form of surgery for at least 5-7 days. Only thereafter should debridement be considered depending on the severity of the bite.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



CAN INFLICT A NASTY BITE



Southern African Python (*Python natalensis*)

Maximum Length:
6 m



Other names:

Luislang, previously called a Rock Python

The Southern African Python is by far our largest snake, reaching a maximum length of around 6 m and weighing in excess of 65 kg. It feeds largely on warm-blooded prey like small antelope, monkeys, game birds and dassies, but also takes leguaans and even crocodiles. Attacks on humans are rare and fatalities virtually unheard of. This snake is largely active at night but is fond of basking during the day. It is at home in water and can remain submerged for long periods. Females produce 30 – 60 (but in exceptional cases more than 100) eggs, which are roughly the size of a tennis ball. The female remains with her eggs throughout incubation and the young measure 50 – 70 cm in length.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Southern African Python (*Python natalensis*)

NON-VENOMOUS, BUT CAN INFLICT A NASTY BITE

Average length: 3 m. Maximum length: 5.5 m

Often confused with: Puff Adders



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



SNAKEBITE PROFILE

SOUTHERN AFRICAN PYTHON

Python natalensis



Risk of Bites

Pythons are ambush hunters and bites usually occur when walking down pathways or through thick bush, where a python is coiled up awaiting a passing prey animal.



Most Bites


Most bites occur to the lower leg, when the snake lunges at a passerby. They are usually quick to release the bitten person, and often leave a row of small teeth punctures in the leg or calf.



Symptoms & Venom

Pythons have no venom, but have over 80 long recurved teeth, which they use to hang onto prey items. Large pythons may leave nasty wounds after a bite.

See pages 103-105
– *Snakes and Snakebite
in Southern Africa*
(2024).




First Aid

Immobilise and reassure the patient. Bandage the bitten area to reduce blood loss and transport the victim to a hospital or doctor.



Medical Treatment

In severe bites victims may require stitches. Bites from pythons should be cleaned and disinfected and monitored for signs of infection. Doctors will also administer tetanus injection.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




MILDLY VENOMOUS



Olive Grass Snake (*Psammophis mossambicus*)

Maximum Length:
1.8 m



Other names:

Olive Whip/sand Snake, Olyf Grasslang

A large, fast-moving diurnal snake that reaches a maximum length of 1.8 m. It is very quick to escape, invariably disappearing down the closest hole but bites readily if handled. Because of its size and habit of lifting its head well off the ground, this snake is often mistaken for a Black Mamba or female Boomslang. It climbs well and is often found in shrubs. Over much of its range, especially northern KZN, it is one of the most frequently encountered snakes. It is often killed on roads. It preys on lizards, frogs, small mammals, birds and other snakes, including venomous ones. The venom of the Olive Grass Snake is mild and has little effect on humans.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Olive Grass Snake (*Psammophis mossambicus*)

MILDLY VENOMOUS

Average length: 80 cm. Maximum length: 1.8 m

Often confused with: Short-snouted Grass Snakes, Black Mambas, Female Boomslang



Distribution is important



Darker spots on upper lip

Olive brown in colour

Tail often has orange tinge

Normally large and fast snakes

Head often has orange tinge

Juveniles may have patterns on neck

Pale belly

www.AFRICANSNAKEBITEINSTITUTE.com



MILDLY VENOMOUS



Western Yellow-bellied Sand Snake (*Psammophis subtaeniatus*)

Maximum Length:
1.4 m



Other names:

Westelike Geelpenssandslang

A fast-moving, slender snake that averages 50-80 cm. This snake is active during the day and is without doubt one of our fastest snakes. It is fond of basking but will move off rapidly if disturbed. It is comfortable in trees and shrubs. Like some of the other snakes in the genus *Psammophis*, it will spin if caught by the tail and snap a piece off. The Western Yellow-bellied Sand snake feeds on lizards, frogs, rodents, small birds and other snakes. The venom of this snake is very mild and has no real effect on humans. Sand snakes, Grass snakes and Skaapstekers can be difficult to tell apart – always check the distribution when attempting to identify one of these snakes.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Western Yellow-bellied Sand Snake (*Psammophis subtaeniatus*)

MILDLY VENOMOUS

Average length: 80 cm - 1 m. Maximum length: 1.4 m

Often confused with: Short-snouted Grass Snakes, Striped Skaapsteker



Longitudinal stripes of
brown, beige and black

Patterns
on neck

Butter-yellow
belly

Black spots
on upper lip

Elongated head

Pale chin

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




MILDLY VENOMOUS



Striped Skaapsteker (*Psammophylax tritaeniatus*)

Maximum Length:
93 cm



Other names:

Gestreepte Skaapsteker

A common diurnal snake that frequents vleis, areas, open grasslands and lowland savannah. It is often found under building rubble near towns. It is very quick to escape and will usually coil around a grass tuft and remain absolutely still. This is one of our fastest-moving snakes. Even when captured it is reluctant to bite. The Skaapstekers and Grass or Sand Snakes are easily confused. Always check the distribution and make sure the snake occurs in your area when attempting to make a positive identification. The Striped Skaapsteker feeds on rodents, frogs, lizards and nestling birds. It has a mild venom that has virtually no effect on humans.



Graham Alexander



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com





ASI EASY ID

Striped Skaapsteker (*Psammophylax tritaeniatus*)

MILDLY VENOMOUS

Average length: 65 cm. Maximum length: 93 cm

Often confused with: Spotted Skaapstekers, and other sand and grass snakes



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




MILDLY VENOMOUS



Eastern Tiger Snake (*Telescopus semiannulatus*)

Maximum Length:
1 m



Other names:
Oostelike Tierslang

A slow-moving nocturnal snake that hunts for lizards, especially geckos, nestling birds, bats and small rodents. It is an excellent climber and spends its days hiding under the bark of trees or under rocks. It can often be seen crossing roads at night. Like the Herald Snake, this snake puts up quite a show when cornered, drawing its head back with coiled neck and striking dramatically with an open mouth. The strikes are rather slow, and the snake may lose its balance. It also flattens the head in the process but does not make a hood. The venom of the Tiger snake is mild and has virtually no effect on humans.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com





ASI EASY ID

Eastern Tiger Snake (*Telescopus semiannulatus*)

MILDLY VENOMOUS

Average length: 40 - 60 cm. Maximum length: 1 m

Often confused with: Cape Coral Snakes, Harlequin Snakes



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



MILDLY VENOMOUS




Photo Luke Kemp

Herald Snake

(Crotaphopeltis hotamboeia)

Maximum Length:

1 m



Other names:

Rooilipslang, Red-lipped Snake

A nocturnal snake that frequents suburban gardens, especially those with water features. It is a toad specialist and often eats toads that look far too large to swallow. The Herald Snake is usually greyish to olive-grey in colour, but some individuals are quite dark. The head is always darker than the body and there may be white speckles on the body. In parts of the range the upper lip is bright red, orange or yellowish to white. When threatened, this snake puts up quite a show, drawing the head back into a striking position with the mouth agape and the head flattened. It is mildly venomous and poses no threat to humans or domestic animals.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Herald Snake (*Crotaphopeltis hotamboeia*)

MILDLY VENOMOUS

Average length: 40 - 60 cm. Maximum length: 82 cm

Can be confused with: juvenile mambas and cobras



Head is distinctly darker
in colour than body.

Dark temples with
iridescent sheen

Dark temples

Upper lips can be red,
orange or white

Body usually light brown,
grey to olive grey

White speckles down the body

Belly is usually white,
to mother of pearl cream

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




CAN INFLICT A NASTY BITE



Mole Snake (*Pseudaspis cana*)

Maximum Length:
2.1 m



Other names:
Molslang

The Mole snake is a muscular day-active snake that spends much of its time down rodent burrows in search of food. It is widespread and occurs throughout most of Southern Africa. It is often encountered in the Cape, but not seen commonly elsewhere in the country. Individuals vary from black to dark brown, medium brown, light brown or even brick red, while the juveniles have bright markings. This snake is not venomous but does have pin-sharp teeth and is capable of quite serious bites that may even require stitches. Mole Snakes are often confused with the Cape Cobra and the venomous stiletto snake even though it does not resemble them.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Mole Snake (*Pseudaspis cana*)

NON-VENOMOUS, BUT CAN INFLICT A NASTY BITE

Average length: 1.2 m. Maximum length: 2.1 m

Often confused with: Cape Cobras, and Snouted Cobras.
Juveniles are often confused with egg-eaters, night adders and skaapstekers




Pointed head



Vary in colour from beige, brown, to reddish brown or black

Bulky snake



Shiny rounded scales



Juvenile Mole Snakes are usually light reddish to greyish brown in colour, with dark, usually zig-zag markings, light spots and mottling down the back

www.AFRICANSNAKEBITEINSTITUTE.com



HARMLESS



Brown House Snake

(*Boaedon capensis*)

Maximum Length:
1.5 m



Other names:

Bruin Huislang

The Brown House Snake, true to its name, frequents human dwellings where it feeds largely on rodents and lizards. It is completely harmless and relies on muscle power to constrict its prey. It is widespread in South Africa and very common in suburban gardens.

This snake is active at night and is relatively slow-moving. Most individuals are medium to dark brown with a mother-of-pearl belly and two light lines on either side of the head – one from the nose through the top of the eye to the back of the head and the other from the back of the eye to the angle of the jaw. The Brown House Snake comes in a variety of shades of brown, often with mottling of white along the neck.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Brown House Snake (*Boaedon capensis*)

HARMLESS

Average length: 55 cm. Maximum length: 1.5 m

Can be confused with: Slug-eaters



Varies in colour from light brown to reddish brown, chocolate brown, tan and pink


May have mottled markings down the neck

One light line from the nose through the top of the eye to the back of the head

Belly is generally a pearl colour

A second light line from the back of the eye to the angle of the jaw

Note the lines above and below the eye



www.AFRICANSNAKEBITEINSTITUTE.com



HARMLESS



Spotted Bush Snake (*Philothamnus semivariegatus*)

Maximum Length:
1.3 m



Other names:

Variegated Bush Snake; Gespikkelde Bosslang

An alert, day-active snake, that climbs well and is often found between the walls and ceilings of outbuildings, where it hunts for geckos. Sadly, this harmless snake is often mistaken for a Green Mamba or Boomslang and needlessly killed.

The body is bright green to a copper-green with distinct blackish spots on the front half of the snake, but little or no dark marks on the back half. The belly is yellowish to white and the pupil orange.

This snake has keeled belly scales which enables it to climb up the bark of a tree or even up face-brick walls. It is commonly found in suburban gardens and juveniles are often killed by domestic cats.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Spotted Bush Snake (*Philothamnus semivariegatus*)

HARMLESS

Average length: 70 cm. Maximum length: 1.3 m

Often confused with: Western Natal Green Snakes,
Green Mambas, Boomslang



Prominent
yellow to red iris


Blackish spots on the
first half of the body

Have bright metallic blue flecks

Excellent climbers, often found in trees and bushes
and in ceilings of outbuildings

In the north of their range,
they often have a
copper coloured tail

Yellowish to white belly
with keeled ventral
and subcaudal scales
for climbing



Double temporal
scales

Three upper
labials making
contact with eye



www.AFRICANSNAKEBITEINSTITUTE.com




HARMLESS



Common Egg-eater (*Dasypeltis scabra*)

Maximum Length:
1.16 m



Other names:

Rhombic Egg-eater, Gewone Eiertvreter

A common snake throughout most of southern Africa except for true desert and closed canopy forests. It averages around 50 cm and is well adapted at climbing trees. It is most abundant in thornveld and grasslands where it can be found in any situation. It is nocturnal, spending most of the day hiding beneath rocks or under loose bark. When agitated, it will coil and uncoil while striking out viciously. The rubbing scales create a rasping sound similar to a hiss. It is completely harmless, and the teeth are reduced. It feeds exclusively on birds' eggs. It may be confused with the venomous Common Night Adder.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com





ASI EASY ID

Common Egg-eater (*Dasypeltis scabra*)

HARMLESS

Average length: 55 cm. Maximum length: 1.16 m

Often confused with: Common Night Adders




www.AFRICANSNAKEBITEINSTITUTE.com

IN THE EVENT OF A SNAKEBITE, DO NOT...

- ✦ Do not try to cut or suck out the venom. Never cut the bite site as cutting may do far more harm than good. Sucking does not remove significant amounts of venom.
- ✦ Do not apply any electric shock therapy. Shock treatments have been tested extensively and have been shown to have no effect on the outcome of the bite. Electric shocks do not neutralise snake venom.
- ✦ Do not apply a tourniquet. Many people are bitten by harmless snakes or venomous snakes that do not inject sufficient venom to cause serious damage. Arterial or venous tourniquets are not advised in most bites as venom is transported largely through the lymphatic system and not through veins. There is little evidence that a tourniquet could be life-saving following a snakebite.
- ✦ Do not apply heat, ice, lotions or potions. Apart from cleaning the bite site with water and applying a sterile gauze dressing nothing should be applied to the bite. Very hot water does not neutralise snake venom.
- ✦ Do not inject antivenom as a first aid measure. Most snakebite victims do not require antivenom and the polyvalent antivenom is only effective for the bites of certain potentially deadly snakes. If required, antivenom must be injected intravenously by a medical doctor and usually in large quantities. As victims may go into anaphylactic shock when injected with antivenom, adrenaline and other drugs may also be required. Anaphylactic shock is a serious medical condition which may be fatal if not treated promptly.

FIRST AID KITS

We stock a wide range of First Aid for Snakebite Kits, as well as Smart Pressure Bandages, Resuscitators, and first aid books. View the range online [here](#).



FIRST AID FOR SNAKEBITE



IN THE EVENT OF A SNAKEBITE, DO...

- ✚ Do get the victim to a hospital as soon as possible and in a safe manner. This is by far the most important first aid measure. For medical advice in a snakebite emergency, call the African Snakebite Institute **+27 (0)82 494 2039**.
- ✚ Do keep the victim calm. Often easier said than done as people tend to panic in a snakebite emergency. Many snakebite victims go into shock, experience a drop in blood pressure and may even become nauseous and faint. If possible, do not allow the victim to walk, run, jump etc. Bear in mind that more than 99% of snakebite victims that are hospitalised within a few hours survive their snakebites.
- ✚ Do remove jewellery as well as tight clothing. If bitten on the hand/arm/foot/lower leg, remove rings, bangles, bracelets, watches, anklets and any other tight jewellery including tight clothing.
- ✚ Do administer artificial respiration if patient stops breathing. In serious snakebite cases involving snakes with predominantly neurotoxic venom, like the Black Mamba or Cape Cobra, the victim may soon experience difficulty with breathing, even within 20 minutes. This is a very serious medical condition and assisted breathing could be life-saving. Mouth-to-mouth resuscitation may be beneficial in the short term, but the proper use of a Bag Valve Mask is far better in such an instance.

SPITTING SNAKES

- ✚ The most common spitting snake in Southern Africa is the Mozambique Spitting. In the western parts of the region this is replaced by the Black Spitting Cobra and the Zebra Cobra. These snakes spray their venom up to a distance of three metres and do so in self-defence to temporarily blind their attacker so that they can make a quick escape.
- ✚ Although spitting snakes accurately aim for the eyes, the venom diffuses into a spray over a wide area and may get onto your hair, face, arms and chest. Venom in the eyes is very painful and must be flushed or diluted as quickly as possible. The production of tears starts this process, but as a first aid measure, flush the eyes with water or a weak (normal) saline solution. The idea is to flush away any excess venom. Transport the victim to a medical doctor, who will examine the eyes for corneal damage and prescribe antibiotic eye drops to prevent secondary infection. There is no reason to use diluted antivenom – it has no proven benefits and may cause an allergic reaction.
- ✚ If treated quickly, the chance of permanent damage to the eyes is extremely remote.

Snakebite Emergency Numbers

Ambulance: 977

Medical Rescue: 911

MRI Botswana Limited: 992

Medical Air Rescue: 390-1601

African Snakebite Institute: +27 82 494 2039

HOSPITALS AROUND BOTSWANA

Princess Marina Hospital
Tel: +267 362 1400
-24.656944, 25.923889

Scottish Livingstone Hospital
Tel: +267 590 8000
-24.41208, 25.48680

Sir Ketumile Masire Teaching Hospital
Tel: +267 355 2048

Sbrana Psychiatric Hospital
Tel: +267 530 5500
-25.199674, 25.678


Nyangabgwe Referral Hospital
Tel: +267 241 3416 / +267 241 1000

Sekgoma Memorial Hospital
Tel: 0800 600 890

Snake Removers Contacts

Find SNAKE REMOVERS in
Botswana
[**HERE**](#)

DOWNLOAD THE
FREE
ASI SNAKES APP
[**HERE**](#)



ASI VEHICLE SPONSORS



The ASI team travels thousands of kilometres a month, often to remote parts of southern Africa, for both snake awareness and venomous snake handling training and for reptile surveys.




We're honoured to have great sponsors, who've made significant changes to our fleet of hardworking vehicles, ensuring they meet our every need.

We'd like to thank our sponsors for making the smooth travel possible.



ASI VEHICLE SPONSORS



MORE RESOURCES



SHOP OUR RANGE OF BOOKS ONLINE [HERE.](#)



CONTACT US

Snake Courses: +27 64 704 7229 or courses@asiorg.co.za

Snake Handling Equipment: +27 60 957 2713 or info@asiorg.co.za

Snake Identification: +27 72 979 1249 or support@asiorg.co.za

Snakebite Emergencies: +27 82 494 2039

WWW.AFRICANSNAKEBITEINSTITUTE.com