



ASI SNAKE BOOKLET

COMMON SNAKES OF BREDASDORP

Copyright: Johan Marais - African Snakebite Institute
First Edition published 2025

814 Wapadrand Rd,
Wapadrand Security Village,
Pretoria, 0081

Cell: +27 (0)82 494 2039
E-mail johan@asiorg.co.za
www.africansnakebiteinstitute.com

All rights reserved. No part of this publication may be reproduced or used commercially without the prior written permission of the copyright owner.

SNAKE COURSES

The ASI offers both public and corporate snake courses.

Public courses take place at various venues throughout South Africa.

The corporate training is done on-site, both in South Africa, as well as in 19 African countries.

We pride ourselves on teaching only the safest snake handling methods, to ensure the safety of both the handler and the snake.

We offer a range of different course options, including snake awareness, first aid for snakebite, venomous snake handling, venomous snake handling bootcamps, as well as advanced first aid for snakebite, advanced venomous snake handling and a reptile photography course.

For course info,
click [here](#) or email us on
courses@asiorg.co.za
for details.



www.AFRICANSNAKEBITEINSTITUTE.com



INTRODUCTION

The African Snakebite Institute is the leading training provider of Snake Awareness, First Aid for Snakebite and Venomous Snake Handling courses in Africa as well as the largest distributor of quality snake handling equipment, snake PPE and snakebite first aid kits. Our aim is to spread snake awareness as far and wide as possible, which is why we've launched our new free E-Book series.

Our CEO, Johan Marais is a well-known herpetologist with over 40 years of experience with snakes. He has worked at various reptile facilities and has collaborated on a variety of scientific studies with some of the world's leading herpetologists. He is also one of Africa's most prominent authors of snake books and has written the best-sellers *A Complete Guide to Snakes of Southern Africa*, *Snakes and Snakebite in Southern Africa* and *First Aid for Snakebite*, to mention just a few.



COMMON SNAKES OF BREDASDORP

The Western Cape has 40 different snake species. Of the 20 snakes that occur around Bredasdorp four are highly venomous, three are venomous and one species can inflict a painful bite. Most snakes are mildly venomous or harmless. We've listed the common species in this booklet, showing venomosity, key ID features, full distribution, and snakebite info for venomous species.

Snakebite is problematic with most bites reported from the hot, wet regions in the eastern part of the country. More than 4,000 snakebites are reported annually with over 1,000 victims being hospitalised. Of those around 60% show signs of envenomation and the remaining 40% show no symptoms – these are referred to as dry bites or may be bites from mildly venomous or harmless snakes. In a recent 5 year survey of 879 snakebite cases in Zululand, nearly 98% of victims showed typical signs of cytotoxic envenomation – largely Mozambique Spitting Cobra, Puff Adder, Common Night Adder and Stiletto Snake bites. Despite their relative frequency, if treated correctly in hospital, these bites are rarely fatal. Most bites occurred in the early evening in the warm wet summer months from November to April/May. Around 84% of the bites were inflicted well below the knee. Less than 11% of these victims received antivenom and none of the bites in this survey were fatal.

Generally speaking we have 10 – 12 fatal snakebites a year in South Africa and these are invariably caused by the Black Mamba and Cape Cobra. Both of these snakes have a potent neurotoxic venom that may quickly affect breathing.

SNAKE HANDLING EQUIPMENT

The African Snakebite Institute is the leading provider of quality snake handling equipment in Africa. View our range online [here](http://www.AFRICANSNAKEBITEINSTITUTE.com).



COMMON SNAKES OF BREDASDORP



Bredasdorp has around twenty different types of snakes. Four species are considered highly venomous while another three can give a bad bite. Most snakes encountered are mildly venomous or harmless to humans.

HIGHLY VENOMOUS

Has caused human fatalities

VENOMOUS

Painful bite, but does not require antivenom

MILDLY VENOMOUS

Not thought to be harmful to humans

HARMLESS

Not dangerous to humans



Cape Cobra
(*Naja nigra*)

HIGHLY VENOMOUS



Rinkhals
(*Hemachatus haemachatus*)

HIGHLY VENOMOUS



Boomslang - male
(*Dispholidus typus*)

HIGHLY VENOMOUS



Puff Adder
(*Bitis arietans*)

HIGHLY VENOMOUS



Herald Snake
(*Crotaphopeltis hotamboeia*)

MILDLY VENOMOUS



Cross-marked Grass Snake
(*Psammophis crucifer*) - Photo Luke Kemp

MILDLY VENOMOUS



Spotted Skaapsteker
(*Psammophylax rhombrostris*)

MILDLY VENOMOUS



Spotted Harlequin Snake
(*Hemiaselaps lacteus*)

MILDLY VENOMOUS



Mole Snake
(*Pseudaspis cana*)

CAN INFLICT A PAINFUL BITE



Mole Snake - juvenile
(*Pseudaspis cana*)

HARMLESS



Brown House Snake
(*Boodon capensis*) - Photo Luke Kemp

HARMLESS



Brown Water Snake
(*Lycodonomorphus rufulus*)

HARMLESS



Olive Snake
(*Lycodonomorphus inornatus*)

HARMLESS



Common Egg-eater
(*Dasypeltis scabra*)

HARMLESS



Common Slug-eater
(*Duberria lutrix*) - Photo Luke Kemp


HARMLESS



Delalande's Beaked Blind Snake
(*Rhinocephalus lalandei*) - Photo Luke Kemp

HARMLESS

JOHAN MARAIS is the author of various books on reptiles including the best-seller *A Complete Guide to Snakes of Southern Africa*. He is a popular public speaker and CEO of the African Snakebite Institute. The African Snakebite Institute (ASI) offers a variety of courses including Snake Awareness, Scorpion Awareness and Venomous Snake Handling. ASI is accredited by the International Society of Zoological Sciences (ISZS) and Travel Doctor-approved service provider. The courses are also accredited by the Health Professions Council of South Africa (HPCSA). ASI is the largest provider of quality snake handling and safety equipment in Africa.



FREE ASI SNAKES APP

Johan Marais | African Snakebite Institute

+27 82 494 2039 | johan@asiorg.co.za

www.AFRICANSNAKEBITEINSTITUTE.com

EMERGENCY PROTOCOL

IN THE EVENT OF A SNAKE BITE

- 1 Keep the victim calm, immobilized and transport the victim to the closest hospital **without delay**.
- 2 If the victim stops breathing, resort to artificial respiration or make use of a Bag Valve Mask.
- 3 Call the Poison Information Centre helpline for further advice: 0861 555 777.

DO NOT

- ... apply a tourniquet.
- ... cut and suck the wound.
- ... use ice or very hot water.
- ... give the victim alcohol.
- ... apply electric shock.
- ... inject antivenom randomly. Antivenom (if required) must be administered by a doctor in a hospital environment.

Download the high resolution poster [here](https://www.asiorg.co.za).

www.AFRICANSNAKEBITEINSTITUTE.com




HIGHLY VENOMOUS



Cape Cobra

(*Naja nivea*)

Maximum Length:
2.3 m



Other names:

Geelslang, Kaapse Kobra, Koper kapel

This snake varies in colour from near black to dark or light brown, beige, yellow or speckled while juveniles have a dark band on the throat. The Cape Cobra is easily confused with the Mole Snake and the Black Spitting Cobra. It is by far our most dangerous cobra and with the Black Mamba it accounts for the majority of fatal snake bites in South Africa. The Cape Cobra may stand its ground if threatened and is quick to form a hood and bite. It occurs largely in the Cape provinces but extends into the Free State, Northwest, southern Botswana and Namibia. The venom is potently neurotoxic causing progressive weakness and may affect breathing rapidly. Victims need urgent hospitalisation and in a severe bite antivenom is essential.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Cape Cobra (*Naja nivea*)

HIGHLY VENOMOUS

Average length: 1.2 - 1.6 m. Maximum length: 2.3 m

Can be confused with: Mole Snakes, Black Spitting Cobras, Snouted Cobras




Often has a dark mark under the eye

Chunky cheeks for venom storage

Varies in colour from light yellow to brown, gold and black.
Often a speckled mix of colours

Spreads an impressive hood

Juveniles have a dark band on throat



Ventral scales often visible on sides

Belly usually same colour as the rest of snake

Usually have dark tipped tails

www.AFRICANSNAKEBITEINSTITUTE.com

Snakebite Profile

Cape Cobra

Naja nivea



Risk of Bites

The Cape Cobra is commonly encountered in suburbia, especially in areas where rodents are abundant. If cornered, it is quick to rear up and form a hood and if approached closely will bite readily.



Most Bites

Most victims are bitten on the lower leg or on a hand, often when attempting to capture or kill the snake.




Symptoms & Venom

A potent and fast-acting neurotoxic venom that is quite similar to Black Mamba venom in its effects. It results in pins and needles in the lips, nausea, excessive sweating, drooping eyelids, difficulty in swallowing and eventually drowsiness and muscle pain followed by paralysis and respiratory failure. Some victims also experience some swelling. In severe cases victims become unresponsive within less than one hour.

See pages 49-51

– *Snakes and Snakebite in Southern Africa* (2024).




First Aid

Immobilise and reassure the patient, who must lie down in a rescue position. Immediately arrange transport to a hospital with emergency facilities. If far from a hospital, apply a pressure bandage (Page 19 of *Snakes and Snakebite in Southern Africa*). Resort to artificial respiration if the victim stops breathing.



Medical Treatment

In cases of severe envenomation patients are intubated and ventilated if they are not breathing. They are treated with 12-15 vials of SAVP Polyvalent antivenom. Additional vials of antivenom may be required at a later stage. Even after receiving antivenom, patients may remain unresponsive for a day or two.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




HIGHLY VENOMOUS



Rinkhals (*Hemachatus haemachatus*)

Maximum Length:
1.5 m



Other names:

Ring-necked Spitting Cobra

The Rinkhals is endemic to Southern Africa and is found only in South Africa, eSwatini, Lesotho and eastern Zimbabwe. Though it resembles a cobra (spreads a hood) it is not a true cobra and gives birth to live young. It is essentially a grassland inhabitant but is also found in fynbos in the Western Cape. It is fond of wetlands where it feeds on frogs. When threatened it is quick to disappear down a hole, but if cornered it will stand its ground, form a hood and spit. The Rinkhals is also quick to play dead with the body turned upside down and the mouth hanging open. The venom of this snake is largely cytotoxic causing pain, swelling and potentially tissue damage. Bites are extremely rare and fatalities virtually unheard of. Polyvalent antivenom is effective.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Rinkhals (*Hemachatus haemachatus*)


HIGHLY VENOMOUS

Average length: 80 cm - 1 m. Maximum length: 1.5 m

Can be confused with: Mozambique Spitting Cobra, Mole Snakes



Adults have black faces.
Juveniles often have
grey faces
with black edged scales



Pointed face

Can spit effectively up to 3 m

Varies in colour from olive to
dark brown, grey or dull black.
In KZN, the Eastern and Western
Cape and Zimbabwe
they are usually banded

Spreads a hood

White bars
on neck region

Has keeled scales,
giving the snake a
rough texture

Belly usually black,
sometimes with grey
or pale edges

Well known for shamming death
when harassed

www.AFRICANSNAKEBITEINSTITUTE.com



SNAKEBITE PROFILE

RINKHALS

Hemachatus haemachatus



Risk of Bites

This snake is quick to hood in a defensive position when startled and will often spit venom at attackers. It is also well known for playing dead when cornered. Bites to humans are uncommon. However, dogs that attack the snake are often bitten and die quickly unless treated with antivenom by a veterinarian soon after a bite.





Most Bites

Bites are not common, except for incidents where snake keepers are bitten by captive snakes. Persons handling the snake may also be bitten.




Symptoms & Venom

The venom is both cytotoxic and neurotoxic. Symptoms include painful swelling of the affected limb and possibly necrosis at the site of the bite. Other symptoms, though not common, include nausea, dizziness and respiratory distress.



See pages 67-69

– *Snakes and Snakebite in Southern Africa* (2024).



First Aid

Immobilise and reassure the patient. Elevate the affected limb. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages. For venom in the eyes, wash the eyes out immediately using water or any other bland liquid and transport the victim to a medical doctor.



Medical Treatment

In cases of severe envenomation where there are clear signs of progressive swelling, which extends more than 10-15cm per hour above the bite site, or respiratory distress, doctors will treat such bites with 12 ampoules (or more) of polyvalent antivenom. Anti-inflammatories (NSAID's) and antibiotics are not indicated in the short term. Fasciotomies are rarely justified and should generally be avoided. In cases where patients suffer tissue damage, surgery may be required but rarely in the first few days. There are no documented fatalities from these bites in more than 45 years.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



HIGHLY VENOMOUS



Boomslang (*Dispholidus typus*)

Maximum Length:
2.1 m



Other names:

Boomslang

The Boomslang is largely tree-living but may descend to the ground to bask. In trees it poses no threat to humans as it is extremely reluctant to bite, and bites are rare. Juveniles are grey with a massive emerald, green eye but change to the adult colours around a length of 1 m. Most males are bright green, sometimes with black between the scales, but in the Cape Provinces they are usually black above with green, yellow or orange sides. Females are usually brown to grey in colour. Boomslang venom is haemotoxic and compromises the blood clotting mechanism, causing uncontrolled bleeding if not treated. The South African Vaccine Producers manufacture a monovalent antivenom for bites from Boomslang that is very effective.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Boomslang (*Dispholidus typus*)


HIGHLY VENOMOUS

Average length: 1.2 m. Maximum length: 2.1 m

Often confused with: Black Mambas, Green Mambas, green snakes and sand snakes




Females are normally olive grey to brown or brick red in colour



Small head with large eye


Scales are keeled giving snake a rough texture



Most males are green in colour



Males in the Cape are usually black above with yellow, green or orange sides



Large eye

Short head



Juveniles have big emerald green eyes

www.AFRICANSNAKEBITEINSTITUTE.com



Snakebite Profile

BOOMSLANG

Dispholidus typus



Risk of Bites

Bites are not common in South Africa. This snake lives in trees and shrubs and interactions with humans are infrequent. It is a placid snake that only bites when provoked or handled. If spotted, it is quick to flee higher up into trees.



Most Bites

Bites occur when a basking snake is accidentally stepped on or grabbed, or when it is handled.




Symptoms & Venom

Drop for drop the Boomslang has the most potent venom of any snake in Africa. It is slow-acting, but potently haemotoxic and symptoms include bleeding from the bite site, the nose, mucous membranes and internal bleeding.

See pages 79-81

– *Snakes and Snakebite in Southern Africa* (2024).




First Aid

Immobilise and reassure the patient. Elevate the affected limb. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages.



Medical Treatment

In cases of severe envenomation patients are treated with monovalent Boomslang antivenom. The initial dosage is two vials but, in some cases, a third vial may be required. This antivenom is produced by the South African Vaccine Producers. It is a Schedule 4 drug, therefore a script is required should one wish to purchase it. It comes in a 10 ml vial and must be refrigerated. It has a three year shelf life.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



HIGHLY VENOMOUS



Puff Adder

(*Bitis arietans*)

Maximum Length:
1.4 m



Other names:

Pofadder, Imbululu

The Puff Adder is widespread over much of southern Africa and further north into Africa. It is a heavy-bodied snake that relies on its excellent camouflage and is reluctant to move. Much of its life is spent in camouflage mode and recent research has shown that when hiding, the Puff Adder will not hiss or strike when approached, as this would give away its presence. It is an ambush hunter that will coil up into a striking position and await its prey. This can last several days or even weeks. Toads are lured closer with flicking of the tongue. Puff Adder venom is potently cytotoxic, causing severe pain, swelling, blistering and in many cases severe tissue damage. Polyvalent antivenom is effective.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Puff Adder (*Bitis arietans*)

HIGHLY VENOMOUS

Average length: 90 cm. Maximum length: 1.4 m

Often confused with: night adders and Berg Adders



www.AFRICANSNAKEBITEINSTITUTE.com



SNAKEBITE PROFILE

PUFF ADDER

Bitis arietans



Risk Of Bites

Bites are common as this snake is widespread throughout most of southern Africa. Along with the Mozambique Spitting Cobra, these two snakes account for the majority of serious snakebites in southern Africa. Dogs are at risk and often get bitten when attempting to kill the snake.



Most Bites

Most victims are bitten at night when accidentally stepping on the snake, but bites are also inflicted during the day when the snake is basking. Most bites are on the feet or lower legs.




Symptoms & Venom

The venom is potently cytotoxic as well as haemotoxic. Symptoms include severe pain, progressive swelling, bleeding from the fang punctures, blistering and tissue damage. The lymph nodes may also be affected. Dogs may perish after a bite from this snake, unless treated with polyvalent antivenom, if required.

See pages 26-28

– *Snakes and Snakebite
in Southern Africa*
(2024).



First Aid


Immobilise and reassure the patient. Elevate the affected limb and remove tight clothing and jewelry. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages.



Medical Treatment

In cases of severe envenomation where there are clear signs of progressive swelling, which extends more than 10-15cm per hour above the bite site, doctors will treat such bites with 6 ampoules (or more) of polyvalent antivenom. Anti-inflammatories (NSAID's) and antibiotics are not indicated in the short term. Fasciotomies are rarely justified and should generally be avoided. In cases where patients suffer tissue damage, surgery may be required but rarely in the first few days.

Fatalities are rare.



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



MILDLY VENOMOUS




Photo Luke Kemp

Herald Snake

(*Crotaphopeltis hotamboeia*)

Maximum Length:
1 m



Other names:

Rooilipslang, Red-lipped Snake

A nocturnal snake that frequents suburban gardens, especially those with water features. It is a toad specialist and often eats toads that look far too large to swallow. The Herald Snake is usually greyish to olive-grey in colour, but some individuals are quite dark. The head is always darker than the body and there may be white speckles on the body. In parts of the range the upper lip is bright red, orange or yellowish to white. When threatened, this snake puts up quite a show, drawing the head back into a striking position with the mouth agape and the head flattened. It is mildly venomous and poses no threat to humans or domestic animals.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Herald Snake (*Crotaphopeltis hotamboeia*)

MILDLY VENOMOUS

Average length: 40 - 60 cm. Maximum length: 82 cm

Can be confused with: juvenile mambas and cobras



Head is distinctly darker
in colour than body.

Dark temples with
iridescent sheen

Dark temples

Upper lips can be red,
orange or white

Body usually light brown,
grey to olive grey

White speckles down the body

Belly is usually white,
to mother of pearl cream

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




MILDLY VENOMOUS



Cross-marked Grass Snake (*Psammophis crucifer*)

Maximum Length:
82 cm



Other names:

Kruismerk Grasslang

One of the smaller grass snakes averaging 30-50 cm but exceeding 82 cm in length. It is active in the day and feeds on small lizards, especially skinks, as well as frogs. It inhabits lowland forest, moist savanna, montane forest, grassland and fynbos. It is common in the sandy scrub-covered areas of the Cape coast and mountain plateau of Mpumalanga. When disturbed it may roll into a tight coil with the head inside, like the Slug-eater. Most individuals are striped with distinct patterns on the side of the head, but some are plain brown to reddish brown with virtually no markings. A mildly venomous snake that poses no threat to humans.



Graham Alexander



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com





ASI EASY ID

Cross-marked Grass Snake (*Psammophis crucifer*)

MILDLY VENOMOUS

Average length: 40 - 60 cm. Maximum length: 82 cm

Often confused with: other sand and grass snakes



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




MILDLY VENOMOUS



Spotted Skaapsteker (*Psammophylax rhombeatus*)

Maximum Length:
1.4 m



Other names:

Rhombic Skaapsteker; Gespikkelde Skaapsteker

A fast-moving diurnal snake that inhabits grasslands, fynbos, moist savanna and semi-desert areas. It is a nervous snake and is quick to escape, either into a bush or grass tuft or under a rock. The name 'Skaapsteker' came from the mistaken belief that these abundant snakes accounted for sheep deaths, when the actual culprit was probably the Cape Cobra. This is also one of few African snakes that coil around and protect their eggs – the Southern African Python being the other. The Spotted Skaapsteker feeds on rodents, lizards, birds, frogs and other snakes. The venom of this snake is mild and poses no threat to humans.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Spotted Skaapsteker (*Psammophylax rhombeatus*)

MILDLY VENOMOUS


Average length: 40 - 60 cm. Maximum length: 1.4 m

Often confused with: other sand and grass snakes and Striped Skaapsteker



Dark patch behind eye

Grassland form




Belly white, with black half bars

Spots on the neck

Cape form



Cape form has red and blue flecks



Striped form

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




MILDLY VENOMOUS



Spotted Harlequin Snake (*Homoroselaps lacteus*)


Maximum Length:
65 cm



Other names:

Gevlekte Kousbandjie

This colourful little snake averages 30 – 40 cm with a maximum length of 65 cm. It varies greatly in colour, depending on locality, and is found in fynbos, lowland forest, moist savanna and grassland. The Spotted Harlequin Snake is quite common in some areas and is found in deserted termite mounds, under rocks and logs or building debris. When exposed it will wriggle violently to escape. It feeds on legless skinks, blind snakes, other snakes and small lizards. Little is known of the venom of this snake, but bites where the snake has been allowed to chew have resulted in pain, swelling and discoloration as well as painful glands. Swelling subsides within 2-3 days.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Spotted Harlequin Snake (*Homoroselaps lacteus*)

MILDLY VENOMOUS

Average length: 40 cm. Maximum length: 65 cm

Often confused with: Cape Coral Snakes and Aurora House Snakes





Grassland form usually black
in colour with yellow spots



Yellow to orange central line
of spots down the length of the snake

Striped form has dark
banding on head and neck

Standard form is a black snake
with yellow transverse bands,
with orange or red infusion



Black eyes

Light splotches
on upper lip

Small head

Dusky yellow belly
with grey infusions

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



CAN INFLICT A NASTY BITE



Mole Snake (*Pseudaspis cana*)

Maximum Length:
2.1 m



Other names:
Molslang

The Mole snake is a muscular day-active snake that spends much of its time down rodent burrows in search of food. It is widespread and occurs throughout most of Southern Africa. It is often encountered in the Cape, but not seen commonly elsewhere in the country. Individuals vary from black to dark brown, medium brown, light brown or even brick red, while the juveniles have bright markings. This snake is not venomous but does have pin-sharp teeth and is capable of quite serious bites that may even require stitches. Mole Snakes are often confused with the Cape Cobra and the venomous stiletto snake even though it does not resemble them.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Mole Snake (*Pseudaspis cana*)

NON-VENOMOUS, BUT CAN INFLICT A NASTY BITE

Average length: 1.2 m. Maximum length: 2.1 m


Often confused with: Cape Cobras, and Snouted Cobras.
Juveniles are often confused with egg-eaters, night adders and skaapstekers



Pointed head



Vary in colour from beige, brown, to reddish brown or black



Bulky snake

Shiny rounded scales



Juvenile Mole Snakes are usually light reddish to greyish brown in colour, with dark, usually zig-zag markings, light spots and mottling down the back

www.AFRICANSNAKEBITEINSTITUTE.com



HARMLESS



Brown House Snake

(*Boaedon capensis*)

Maximum Length:
1.5 m



Other names:

Bruin Huislang

The Brown House Snake, true to its name, frequents human dwellings where it feeds largely on rodents and lizards. It is completely harmless and relies on muscle power to constrict its prey. It is widespread in South Africa and very common in suburban gardens.

This snake is active at night and is relatively slow-moving. Most individuals are medium to dark brown with a mother-of-pearl belly and two light lines on either side of the head – one from the nose through the top of the eye to the back of the head and the other from the back of the eye to the angle of the jaw. The Brown House Snake comes in a variety of shades of brown, often with mottling of white along the neck.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Brown House Snake (*Boaedon capensis*)

HARMLESS

Average length: 55 cm. Maximum length: 1.5 m

Can be confused with: Slug-eaters



Varies in colour from light brown to reddish brown, chocolate brown, tan and pink

May have mottled markings down the neck

One light line from the nose through the top of the eye to the back of the head

Belly is generally a pearl colour

A second light line from the back of the eye to the angle of the jaw

Note the lines above and below the eye



www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



HARMLESS



Brown Water Snake (*Lycodonomorphus rufulus*)

Maximum Length:
97 cm



Other names:
Bruin Waterslang

A smallish snake averaging 30-50 cm with a maximum length of 97 cm. It is light to dark brown with a salmon-pink to yellow coloured belly. Snakes from the Western Cape are often much darker. This snake is not as aquatic as the Dusky-bellied Water Snake but does frequent rivers, streams, vleis and other damp areas. It is active at night and swims well but is often seen hunting along shaded streams during the day. It is a powerful constrictor and feeds on frogs, tadpoles, fish and occasionally nestling rodents and geckos. Females produce 6 – 23 eggs in summer. Though harmless this snake is much feared in Zulu culture and is called Ivuzamanzi elimdubu.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Brown Water Snake (*Lycodonomorphus rufulus*)

HARMLESS

Average length: 45-60 cm. Maximum length: 97 cm

Often confused with: Olive Snakes, Yellow-bellied House Snakes



www.AFRICANSNAKEBITEINSTITUTE.com




HARMLESS



Olive Snake

(Lycodonomorphus inornatus)

Maximum Length:
1.3 m



Other names:

Olive House/Ground Snake, Olyfslang

A slow-moving nocturnal snake that is quite similar to the Brown House Snake in behaviour. It averages 30-60 cm but reaches 1.3 m in length. Commonly found around houses but prefers damp areas where it hunts for lizards, rodents and other snakes. Because of its colour, varying from olive-green to light brown or black, it is mistaken for a variety of other snakes including the Black Mamba and the Brown Water Snake.

The Olive Snake used to be in the same genus as the house snakes, hence the common name Olive House Snake, but has since been moved to the same genus as the water snakes. This snake is not venomous and poses no threat to humans.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com



ASI EASY ID

Olive Snake (*Lycodonomorphus inornatus*)

HARMLESS

Average length: 75 cm. Maximum length: 1.3 m

Often confused with: Brown House Snakes, Brown Water Snakes, Yellow-Bellied House Snakes



Adults are olive green to brown or black in colour

Juveniles in the Cape are light olive green with distinctly dark head

Juveniles in the East are usually pitch black

Adults often have pale chin

Uniform in colour, including belly

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




HARMLESS



Common Egg-eater (*Dasypeltis scabra*)

Maximum Length:
1.16 m



Other names:

Rhombic Egg-eater, Gewone Eiervreter

A common snake throughout most of southern Africa except for true desert and closed canopy forests. It averages around 50 cm and is well adapted at climbing trees. It is most abundant in thornveld and grasslands where it can be found in any situation. It is nocturnal, spending most of the day hiding beneath rocks or under loose bark. When agitated, it will coil and uncoil while striking out viciously. The rubbing scales create a rasping sound similar to a hiss. It is completely harmless, and the teeth are reduced. It feeds exclusively on birds' eggs. It may be confused with the venomous Common Night Adder.



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com





ASI EASY ID

Common Egg-eater (*Dasypeltis scabra*)

HARMLESS

Average length: 55 cm. Maximum length: 1.16 m

Often confused with: Common Night Adders



www.AFRICANSNAKEBITEINSTITUTE.com



HARMLESS



Common Slug-eater (*Duberria lutrix*)

Maximum Length:
43 cm



Other names:

Gewone Slakvreter, Tabakrolletjie

A small snake that averages 30 – 35 cm. It feeds exclusively on snails and slugs and locates them by following the slime trail. The Common Slug-eater favours damp localities in the south-eastern half of southern Africa.

During the day it seeks shelter under rocks, logs and grass tufts and emerges in the evening to hunt. It is reluctant to bite and if threatened will defecate and secrete a foul-smelling substance simultaneously. It also has the peculiar habit of rolling up into a tight spiral with the head concealed, like a roll of tobacco, hence the Afrikaans name. This small slow-moving snake is not venomous and poses no threat to humans.



Luke Kemp



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Common Slug-eater (*Duberria lutrix*)

HARMLESS

Average length: 35 cm. Maximum length: 45 cm

Often confused with: Juvenile Cape Cobras



www.AFRICANSNAKEBITEINSTITUTE.com




HARMLESS



Delalande's Beaked Blind Snake (*Rhinotyphlops lalandei*)

Maximum Length:
35 cm



Other names:

Delalande se Blindeslang

A widespread burrowing snake that spends most of its life underground in search of ants and their eggs. Individuals may be found in soil under rocks or logs, in deserted termite mounds or on the surface at night, especially after rains. They are pale pink to dark-slate or grey-brown in colour. Each scale is pale-edged giving it a checkered effect. The young are pink or flesh-coloured. It is a small snake with adults averaging around 25 - 30 cm, with a maximum size of about 35 cm. They are easily confused with other blind snakes or worm snakes. Blind snakes are preyed upon by birds and a variety of snakes including the Spotted Harlequin Snake.



Jake Kemp



© Johan Marais African Snakebite Institute 2022

www.AFRICANSNAKEBITEINSTITUTE.com

www.AFRICANSNAKEBITEINSTITUTE.com




ASI EASY ID

Delalande's Beaked Blind Snake (*Rhinotyphlops lalandei*)

HARMLESS

Average length: 30 cm. Maximum length: 38.5 cm


Often confused with: stiletto snakes and Bibron's Blind Snakes



Large rostral scale
which does not
quite extend to the eye

Small eye, covered
by scale

Colour varies from
pink to mauve.
Juveniles are
flesh-coloured



Top is often darker
with chequered effect

Belly pink to grey
or straw-coloured

www.AFRICANSNAKEBITEINSTITUTE.com

FIRST AID FOR SNAKEBITE



IN THE EVENT OF A SNAKEBITE, DO...

- ✦ Do get the victim to a hospital as soon as possible and in a safe manner. This is by far the most important first aid measure. For medical advice in a snakebite emergency, call the African Snakebite Institute **+27 (0)82 494 2039**.
- ✦ Do keep the victim calm. Often easier said than done as people tend to panic in a snakebite emergency. Many snakebite victims go into shock, experience a drop in blood pressure and may even become nauseous and faint. If possible, do not allow the victim to walk, run, jump etc. Bear in mind that more than 99% of snakebite victims that are hospitalised within a few hours survive their snakebites.
- ✦ Do remove jewellery as well as tight clothing. If bitten on the hand/arm/foot/lower leg, remove rings, bangles, bracelets, watches, anklets and any other tight jewellery including tight clothing.
- ✦ Do administer artificial respiration if patient stops breathing. In serious snakebite cases involving snakes with predominantly neurotoxic venom, like the Black Mamba or Cape Cobra, the victim may soon experience difficulty with breathing, even within 20 minutes. This is a very serious medical condition and assisted breathing could be life-saving. Mouth-to-mouth resuscitation may be beneficial in the short term, but the proper use of a Bag Valve Mask is far better in such an instance.

SPITTING SNAKES

- ✦ The common spitting snakes in South Africa are the Mozambique Spitting Cobra or M'Fezi and the Rinkhals. In the western parts of the country these are replaced by the Black Spitting Cobra. These snakes spray their venom up to a distance of three metres and do so in self-defence to temporarily blind their attacker so that they can make a quick escape.
- ✦ Although spitting snakes accurately aim for the eyes, the venom diffuses into a spray over a wide area and may get onto your hair, face, arms and chest. Venom in the eyes is very painful and must be flushed or diluted as quickly as possible. The production of tears starts this process, but as a first aid measure, flush the eyes with water or a weak (normal) saline solution. The idea is to flush away any excess venom. Transport the victim to a medical doctor, who will examine the eyes for corneal damage and prescribe antibiotic eye drops to prevent secondary infection. There is no reason to use diluted antivenom – it has no proven benefits and may cause an allergic reaction.
- ✦ If treated quickly, the chance of permanent damage to the eyes is extremely remote.

Snakebite Emergency Numbers

Netcare Ambulance: 082 911

ER24 Ambulance: 084 124

Provincial Ambulance: 10177/112

African Snakebite Institute: +27 (0)82 494 2039

HOSPITALS AROUND BREDASDORP

Otto du Plessis Hospital

Tel: 028 424 2652

(-34.536718, 20.033912)

Bredasdorp Cnr Dorpsig & Van
Riebeeck Street, Bredasdorp, Cape
Agulhas, 7280

Swellendam Provincial Hospital

Tel: 028 514 1141

(-34.024516, 20.449789)

47 Drostdy st, Swellendam, 6740



Snake Removers Contacts

Find SNAKE REMOVERS in
Bredasdorp

HERE

DOWNLOAD THE
FREE
ASI SNAKES APP
HERE



www.AFRICANSNAKEBITEINSTITUTE.com

ASI VEHICLE SPONSORS



The ASI team travels thousands of kilometres a month, often to remote parts of southern Africa, for both snake awareness and venomous snake handling training and for reptile surveys.




We're honoured to have great sponsors, who've made significant changes to our fleet of hardworking vehicles, ensuring they meet our every need.

We'd like to thank our sponsors for making the smooth travel possible.



ASI VEHICLE SPONSORS



MORE RESOURCES



SHOP OUR RANGE OF BOOKS ONLINE [HERE](#).



CONTACT US

Snake Courses: +27 64 704 7229 or courses@asiorg.co.za

Snake Handling Equipment: +27 60 957 2713 or info@asiorg.co.za

Snake Identification: +27 72 979 1249 or support@asiorg.co.za

Snakebite Emergencies: +27 82 494 2039

WWW.AFRICANSNAKEBITEINSTITUTE.com