

Copyright: Johan Marais - African Snakebite Institute First Edition published 2025

814 Wapadrand Rd, Wapadrand Security Village, Pretoria, 0081

Cell: +27 (0)82 494 2039 E-mail johan@asiorg.co.za www.africansnakebiteinstitute.com

All rights reserved. No part of this publication may be reproduced or used commercially without the prior written permission of the copyright owner.

SNAKE COURSES

The ASI offers both public and corporate snake courses.

Public courses take place at various venues throughout South Africa.

The corporate training is done on-site, both in South Africa, as well as in 19 African countries.

We pride ourselves on teaching only the safest snake handling methods, to ensure the safety of both the handler and the snake.

We offer a range of different course options, including snake awareness, first aid for snakebite, venomous snake handling, venomous snake handling bootcamps, as well as advanced first aid for snakebite, advanced venomous snake handling and a reptile photography course.

For course info, click here or email us on courses@asiorg.co.za for details.





INTRODUCTION

he African Snakebite Institute is the leading training provider of Snake Awareness, First Aid for Snakebite and Venomous Snake Handling courses in Africa as well as the largest distributer of quality snake handling equipment, snake PPE and snakebite first aid kits. Our aim is to spread snake awareness as far and wide as possible, which is why we've launched our new free E-Book series.

ur CEO, Johan Marais is a well-known herpetologist with over 40 years of experience with snakes. He has worked at various reptile facilities and has collaborated on a variety of scientific studies with some of the world's leading herpetologists. He is also one of Africa's most prominent authors of snake books and has written the best-sellers A Complete Guide to Snakes of Southern Africa, Snakes and Snakebite in Southern Africa and First Aid for Snakebite, to mention just a few.



COMMON SNAKES OF ZIMBABWE

Zimbabwe has 72 different snake species. Twelve of these are highly venomous, nine are venomous, two can inflict deep lacerations and the rest are mildly venomous or harmless. We've listed the common species in this booklet, showing venomiosity, key ID features, full distribution, and snakebite info for venomous species.

Snakebite is problematic with most bites reported from the hot wet regions in the eastern part of the country. More than 8,000 snakebites are reported annually with over 1,000 victims being hospitalised. Of those around 60% show signs of envenomation and the remaining 40% show no symptoms – these are referred to as dry bites. In a recent survey of 879 snakebite cases in Zululand over a five year period nearly 98% of victims showed typical signs of cytotoxic envenomation – largely Mozambique Spitting Cobra, Puff Adder, Rhombic Night Adder and Stiletto snake bites. Mozambique Spitting Cobra and Puff Adder bites are rarely fatal.

ost bites occur in the early evening in the warm wet summer months from November to April/May. Around 84% of bites are inflicted well below the knee. Less than 11% of these victims received antivenom and none of the bites in this survey were fatal. Generally speaking we have 10 – 12 fatal snakebites a year in South Africa and these are invariably caused by the Black Mamba and Cape Cobra. Both of these snakes have a potent neurotoxic venom that may quickly affect breathing.

SNAKE HANDLING EQUIPMENT

The African Snakebite Institute is the leading provider of quality snake handling equipment in Africa. View our range online here.



COMMON SNAKES OF ZIMBABW



Zimbabwe has around 72 species of snakes. Twelve species are considered highly venomous. Nine species are venomous and another two can deliver a painful bite that may require stitches. Twenty species are mildly venomous and twenty-nine are harmless to humans.

HIGHLY

human fatalities

VENOMOUS

Painful bite, but does not require antivenom

Not thought to be harmful HARMLESS

Not dangerous to humans



Black Mamba



Snouted Cobra









Puff Adder



Snouted Night Adder





Herald Snake



Eastern Tiger Snake



non Purple-glossed Snake











Common Wolf Snake









JOHAN MARAIS is the author of various books on reptiles including the best-seller A Complete Guide to Snakes of Southern Africa. He is a popular public speaker and CEO of the African Snakebite Institute. The African Snakebite Institute (ASI) offers a variety of courses including Snake Awareness, Scorpion Awareness and Venomous Snake Handling. ASI is accredited by the International Society of Zoological Sciences (ISZS) and Travel Doctor-approved service provider. The courses are also accredited by the Health Professions Council of South Africa (HPCSA). ASI is the largest provider of quality snake handling and safety equipment in Africa.

Johan Marais | African Snakebite Institute

+27 82 494 2039 | johan@asiorg.co.za

www.AFRICANSNAKEBITEINSTITUTE.com

find trained snake removers in

your area, browse our free resources, and more



Download the high resolution poster here.







Black Mamba

(Dendroaspis polylepis)

Maximum Length: 3.8 m



Other names:

Swartmamba


This is by far the largest venomous snake in Africa, with a maximum length of 3.8 m (historically specimens over 4 m were reported). Despite its reputation it is a shy, elusive snake that is quick to escape but will not hesitate to strike repeatedly if cornered. It is active during the day, often basking near a hole in an anthill or large rock crevice, into which it quickly disappears into if disturbed. When threatened it will gape exposing the black inner lining of the mouth and it may form a narrow hood. The venom of this snake is potently neurotoxic and may cause difficulty with breathing within half an hour. Antivenom is effective but often required in large quantities (10 – 15 vials).







© Johan Marais African Snakebite Institute 2022





SNAKEBITE PROFILE **BLACK MAMBA**

Dendroaspis polylepis



Risk of Bites

The Black Mamba is a nervous snake and is quick to escape. If accidentally cornered in a confined space or surprised in the veld, it may strike out and bite more than once in quick succession. Contrary to popular stories, it does not chase people and bites are quite rare.



Most Bites

Most victims are bitten in the veld when they accidentally get very close to a Black Mamba basking. Or when accidentally cornering a mamba in a confined space like a pump house on a farm. Never approach a Black Mamba closely and do not try to kill or catch one.



Symptoms & Venom

A potent and fast-acting neurotoxic venom that results in pins and needles in the lips, hands and feet, as well as nausea, excessive sweating, drooping eyelids, difficulty in swallowing and eventually drowsiness and muscle pain followed by paralysis and respiratory failure. In severe cases victims become unresponsive within less than one

See pages 43-45 - Snakes and Snakebite in Southern Africa (2024).



First Aid

Immobilise and reassure the patient, who must lie down in a rescue position. Immediately arrange transport to a hospital with emergency facilities. If far from a hospital, apply a pressure bandage (Page 19 of Snakes and Snakebite in Southern Africa). Resort to artificial respiration if the victim stops breathing.



Medical Treatment

In cases of severe envenomation where patients are not breathing, they are intubated and ventilated and treated with 12-15 vials of SAVP Polyvalent antivenom. Additional vials of antivenom may be required at a later stage.









Snouted Cobra (Naja annulifera)

Maximum Length: 2.5m



Other names:

Wipneuskobra, Bosveldkapel

The Snouted Cobra is largely a moist and dry savannah inhabitant but is also found in lowveld and bushveld areas where it actively hunts its prey during the day or at night. It feeds on rodents, birds, their eggs and especially toads, but is fond of eating snakes, including the Puff Adder. It is not an aggressive snake but if cornered will stand its ground and form an impressive hood. It may exceed 2.5 m in length. It is usually different shades of brown, but some individuals have broad light brown and purplish-black bands on the back. The venom of this snake is both neurotoxic and cytotoxic and may cause some tissue damage. Polyvalent antivenom is effective, but luckily bites are rare.







© Johan Marais African Snakebite Institute 2022




Snouted Cobra (Naja annulifera)

HIGHLY VENOMOUS

Average length: 1.2 - 1.6 m. Maximum length: 2.5 m

Can be confused with: Mozambique Spitting Cobras, Cape cobras







SNAKEBITE PROFILE SNOUTED COBRA

Naja annulifera



Risk of Bites

Bites from the Snouted Cobra are not common. This snake frequently visits residential gardens in search of rodents and toads. It also raids poultry runs. Dogs are at risk and often get bitten when attempting to kill the snake.



Most Bites

Bites are infrequent and usually occur when accidentally confronted or stepped on at night. Most bites are on the lower legs, or to those who try and handle the snake.



& Symptoms & Venom

The venom is both neurotoxic and cytotoxic. It is one of our largest cobras reaching 2,5 m in length and can deliver a lot of venom in a single bite. Bites may result in burning pain, nausea and swelling followed by tissue damage. It can also affect breathing.



See pages 52-53 - Snakes and Snakebite in Southern Africa (2024).



First Aid

Immobilise and reassure the patient, who must lie down in a rescue position. Immediately arrange transport to a hospital with emergency facilities. If far from a hospital, apply a pressure bandage (Page 19 of Snakes and Snakebite in Southern Africa). Resort to artificial respiration if the victim stops breathing.



開 Medical Treatment

In cases of severe envenomation, especially where patients are not breathing, they should be intubated, ventilated, and treated with 12-15 vials of SAVP Polyvalent antivenom. Additional vials of antivenom may be required at a later stage.







HIGHLY VENOMOUS

Mozambique Spitting Cobra (Naja mossambica)

Maximum Length: 1.8 m



Other names:

Mosambiekse Spoegkobra

The Mozambique Spitting Cobra occurs on the eastern side of South Africa, most of Mozambique, Zimbabwe, north-eastern Botswana, northern Namibia and further north in Africa. Its diet consists largely of frogs, small mammals, birds and snakes, including the Puff Adder. It is active on overcast days, but more active at night, often ending up in houses where people are bitten while asleep. This snake accounts for the majority of serious snakebite cases in southern Africa, but fatalities are rare. It also spits its venom and does not always spread a hood when doing so. The venom is potently cytotoxic causing pain, swelling, blisters and in many cases severe tissue damage. Antivenom is effective if administered early and in large dosages.







© Johan Marais African Snakebite Institute 2023




Mozambique Spitting Cobra (Naja mossambica)

HIGHLY VENOMOUS

Average length: 80 cm - 1.2 m. Maximum length: 1.83 m

Can be confused with: Rinkhals, Snouted Cobras





SNAKEBITE PROFILE

MOZAMBIQUE SPITTING COBRA

Naja mossambica



Risk of Bites

Bites are common, as this snake is widespread throughout much of southern Africa. Along with the Puff Adder, these two snakes account for most serious snakebites in southern Africa. Dogs are at risk and often get bitten when attempting to kill the snake.




Most Victims are bitten at night when accidentally stepping on the snake, or while in bed. The snake enters homes and may bite sleeping victims, often multiple times. Bites in bed appear to be somewhat of a feeding response with the snake smelling a mammal and biting. Should you live within the range of this snake it is advisable to ensure that exit doors seal well without gaps or to fix mosquito-proof doors to all exits. People are also encouraged to sleep under mosquito nets.



& Symptoms & Venom

The venom is predominantly and potently cytotoxic. Symptoms include severe pain, progressive swelling, various lesions and, in some cases, severe tissue damage.



See pages 64-65 Snakes and Snakebite in Southern Africa (2024).



First Aid

Immobilise and reassure the patient. Elevate the affected limb and remove tight clothing and jewelry. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages. For venom in the eyes, wash the eyes out immediately using water or any other bland liquid and get the victim to a medical doctor for further examination and treatment.



Medical Treatment

In cases of severe envenomation where there are clear signs of progressive swelling, extending more than 10-15cm per hour above the bite site, doctors must treat such bites with 12 ampoules (or more) of polyvalent antivenom. It appears that SAVP polyvalent antivenom is not very effective against the venom of this snake, therefore it should be administered within an hour or two after a bite. Anti-inflammatories (NSAID's) and antibiotics are not indicated in the short term. Fasciotomies are rarely justified and should generally be avoided. In cases where patients suffer tissue damage, surgery may be required but rarely in the first few days. Fatalities are rare.










Boomslang

(Dispholidus typus)

Maximum Length: 2.1 m



Other names:

Boomslang


The Boomslang is largely tree-living but may descend to the ground to bask. In trees it poses no threat to humans as it is extremely reluctant to bite, and bites are rare. Juveniles are grey with a massive emerald, green eye but change to the adult colours around a length of 1 m. Most males are bright green, sometimes with black between the scales, but in the Cape Provinces they are usually black above with green, yellow or orange sides. Females are usully brown to grey in colour. Boomslang venom is haemotoxic and compromises the blood clotting mechanism, causing uncontrolled bleeding if not treated. The South African Vaccine Producers manufacture a monovalent antivenom for bites from Boomslang that is very effective.







O Johan Marais African Snakebite Institute 2023





SNAKEBITE PROFILE **BOOMSLANG**

Dispholidus typus



Risk of Bites

Bites are not common in South Africa. This snake lives in trees and shrubs and interactions with humans are infrequent. It is a placid snake that only bites when provoked or handled. If spotted, it is quick to flee higher up into trees.



Most Bites


Bites occur when a basking snake is accidentally stepped on or grabbed, or when it is handled.



Symptoms & Venom

Drop for drop the Boomslang has the most potent venom of any snake in Africa. It is slow-acting, but potently haemotoxic and symptoms include bleeding from the bite site, the nose, mucous membranes and internal bleeding.

See pages 79-81 Snakes and Snakebite in Southern Africa (2024).




First Aid

Immobilise and reassure the patient. Elevate the affected limb. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages.



Medical Treatment

In cases of severe envenomation patients are monovalent Boomslang treated with antivenom. The initial dosage is two vials but, in some cases, a third vial may be required. This antivenom is produced by the South African Vaccine Producers. It is a Schedule 4 drug, therefore a script is required should one wish to purchase it. It comes in a 10 ml vial and must be refrigerated. It has a three year shelf life.









Southern Vine Snake (Thelotornis capensis)

Maximum Length: 1.47 m



Other names:

Twig Snake; Bird Snake; Takslang; Voëlslang This perfectly-camouflaged tree-living snake is seldom seen because of its excellent camouflage and habit of remaining very still in low shrubs, observing the ground below for passing lizards and snakes. Birds often mob this snake, and it will inflate its neck with its bright orange tongue flickering – this leads to the incorrect assumption that they lure birds with their tongue. It is extremely placid but, if provoked, will inflate its neck and strike viciously. Most bites are to snake handlers. Like the Boomslang, the venom of this snake is haemotoxic affecting the blood clotting mechanism and causing uncontrolled bleeding. There is no antivenom for the venom of this snake and although a few fatalities have been reported, none were in South Africa.







© Johan Marais African Snakebite Institute 2022




Southern Vine Snake (Thelotornis c. capensis)

HIGHLY VENOMOUS

Average length: 1.2 m. Maximum length: 1.47 m

Often confused with: Juvenile Boomslang, Eastern Bark Snakes







SNAKEBITE PROFILE SOUTHERN VINE SNAKE

Thelotornis capensis



Risk Of Bites

This snake lives in trees and shrubs and interactions with humans are not common. It is a placid snake that only bites when provoked or handled. Bites are very rare in South Africa.



Most Bites

There are very few documented bites from this snake in southern Africa, except for incidents where snake keepers are bitten by captive snakes, or by other people handling this snake.



Symptoms & Venom

The venom is potently haemotoxic, like that of the Boomslang. Symptoms include bleeding from the nose. mucous membranes and bite site. This may eventually lead to internal bleeding.

See pages 82-84 - Snakes and Snakebite in Southern Africa (2024).




First Aid

Immobilise and reassure the patient. Elevate the affected limb. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages.



Medical Treatment

Polyvalent and Monovalent antivenom do not offer cross coverage for Vine Snake bites and should not be administered. Bites are treated symptomatically with transfers and platelets, but this may not be effective in severe bites. To date we do not have any documented fatalities in South Africa, however a few fatal bites have been recorded further north in Africa.






HIGHLY VENOMOUS



Puff Adder

(Bitis arietans)

Maximum Length: 1.4 m



Other names:

Pofadder, Imbululu

The Puff Adder is widespread over much of southern Africa and further north into Africa. It is a heavy-bodied snake that relies on its excellent camouflage and is reluctant to move. Much of its life is spent in camouflage mode and recent research has shown that when hiding, the Puff Adder will not hiss or strike when approached, as this would give away its presence. It is an ambush hunter that will coil up into a striking position and await its prey. This can last several days or even weeks. Toads are lured closer with flicking of the tongue. Puff Adder venom is potently cytotoxic, causing severe pain, swelling, blistering and in many cases severe tissue damage. Polyvalent antivenom is effective.







© Johan Marais African Snakebite Institute 2022




Puff Adder (Bitis arietans)

HIGHLY VENOMOUS

Average length: 90 cm. Maximum length: 1.4 m

Often confused with: night adders and Berg Adders







SNAKEBITE PROFILE PUFF ADDER

Bitis arietans



Risk Of Bites

Bites are common as this snake is widespread throughout most of southern Africa. Along with the Mozambique Spitting Cobra, these two snakes account for the majority of serious snakebites in southern Africa. Dogs are at risk and often get bitten when attempting to kill the snake.



Most Bites


Most victims are bitten at night when accidentally stepping on the snake, but bites are also inflicted during the day when the snake is basking. Most bites are on the feet or lower legs.



Symptoms & Venom

The venom is potently cytotoxic as well as haemotoxic. Symptoms include severe pain, progressive swelling, bleeding from the fang punctures, blistering and tissue damage. The lymph nodes may also be affected. Dogs may perish after a bite from this snake, unless treated with polyvalent antivenom, if required.

See pages 26-28 Snakes and Snakebite in Southern Africa (2024).





Immobilise and reassure the patient. Elevate the affected limb and remove tight clothing and jewelry. Promptly transport the victim to the nearest hospital that has emergency facilities. Avoid all other first aid remedies including bandages.



Medical Treatment

In cases of severe envenomation where there are clear signs of progressive swelling, which extends more than 10-15cm per hour above the bite site, doctors will treat such bites with 6 ampoules (or more) of polyvalent antivenom. Anti-inflammatories (NSAID's) and antibiotics are not indicated in the short term. Fasciotomies are rarely justified and should generally be avoided. In cases where patients suffer tissue damage, surgery may be required but rarely in the first few days.

Fatalities are rare.










VENOMOUS

Snouted Night Adder

(Causus defilippii)

Maximum Length: 43 cm



Other names: Wipneusnagadder


The Snouted Night Adder is very similar to the Common Night Adder in appearance and behaviour, but is much smaller, averaging 35 cm, with a maximum length of 43 cm. This snake can be distinguished from the Common Night Adder by its upturned snout. The dark V on the nape is usually broad and there may be a darker band down the back. Like the Common Night Adder, it feeds almost exclusively on frogs, particularly toads. Snouted Night Adder venom is similar to that of the Common Night Adder – cytotoxic, causing pain and swelling. Because of its size, the venom yield is less than the bigger Common Night Adder and bites are seldom serious. Antivenom is not effective and not required.







O Johan Marais African Snakebite Institute 2023





SNAKEBITE PROFILE **SNOUTED NIGHT ADDER**

Causus defilippii



Risk of Bites

This is a small snake, which is not commonly encountered. Very few bites have been recorded.



Most Bites

Bites mainly occur when stepping on this snake.



Symptoms & Venom

The venom of the Snouted Night Adder is mild, causing pain and swelling. The swelling usually subsides in a few days.

See pages 18-22 - Snakes and Snakebite in Southern Africa (2024).




First Aid

Immobilise and reassure the patient and elevate the affected limb. Remove tight clothing and jewelry. Transport the victim to the nearest hospital or doctor for pain management. Avoid all other first aid remedies including bandages.



Medical Treatment

There is no antivenom for bites from this snake, and victims would be treated








VENOMOUS

Common Night Adder (Causus rhombeatus)

Maximum Length: 1 m



Other names:

Rhombic Night Adder, Gewone Nagadder

Despite the common name, the Night Adder is more active during the day. It averages 30 – 60 cm in length, but can reach 1 m. It can be identified by the distinct dark V-marking on the head. It prefers damp localities and is a frog specialist. The Night Adder is generally a docile snake but will hiss aggressively and strike when provoked. It accounts for many bites, especially in KZN and the Eastern Cape. It is commonly found in suburban gardens.

Night Adder venom is cytotoxic, and most bites are not serious – causing pain and swelling. Having said that, the venom of this snake should never be underestimated, especially in children, as some victims require hospitalisation.







© Johan Marais African Snakebite Institute 2022



Common Night Adder (Causus rhombeatus)

VENOMOUS

Average length: 40 - 60 cm. Maximum length: 1 m

Often confused with: Rhombic Egg-eaters, Berg Adders and Puff Adders



Despite the common name, they are mainly active during the day Single distinct "V" Rhombic markings. marking on the head. Dark brown, light-edged blotches The apex of the V ending along the centre of the body and tail between the eyes Black lines on the Often has edges of the scales dark mask of the upper lip over eye Belly usually white to yellowish or light grey Colour, varying shades of light grey, olive, Round pupil or light to pinkish brown



SNAKEBITE PROFILE **COMMON NIGHT ADDER**

Causus rhombeatus



Risk of Bites

Bites are quite common in coastal KwaZulu-Natal and the Eastern Cape. Despite the common name, this snake is active during the day and frequents moist areas at dams, rivers and streams, where it feeds largely on toads. Dogs are at risk and often get bitten when attempting to kill the snake.



Most Bites

People are bitten when accidentally stepping onto one of these snakes. Most bites are on the feet or lower legs. Some people are under the wrong impression that Night Adder venom is mild and get bitten when handling the snake.



First Aid

Immobilise and reassure the patient, elevate the affected limb and rest it. In severe cases, especially in small children, the patient may require hospitalization for rehydration and pain.

See pages 40-41 Snakes and Snakebite in Southern Africa (2024).





Symptoms & Venom


cytotoxic is largely (cell-destroying), causing pain, swelling, and discoloration of the bitten area within an hour or two. The lymph nodes may also be affected. Swelling and discoloration may persist for more than a week, but blistering is rare. The venom of this snake does not cause tissue damage but could be quite serious when dogs are bitten. Small dogs may perish after a bite from this snake.



Medical Treatment

There is no antivenom for this bite of a Common Night Adder. Both antibiotics and anti-inflammatories (NSAIDs) are not indicated and should not be given in the first 72 hours. If well rested, patients recover fully within a week or two.












Herald Snake

(Crotaphopeltis hotamboeia)

Maximum Length: 1 m



Other names:

Rooilipslang, Red-lipped Snake


A nocturnal snake that frequents suburban gardens, especially those with water features. It is a toad specialist and often eats toads that look far too large to swallow. The Herald Snake is usually greyish to olive-grey in colour, but some individuals are quite dark. The head is always darker than the body and there may be white speckles on the body. In parts of the range the upper lip is bright red, orange or yellowish to white. When threatened, this snake puts up quite a show, drawing the head back into a striking position with the mouth agape and the head flattened. It is mildly venomous and poses no threat to humans or domestic animals.







O Johan Marais African Snakebite Institute 2023








MILDLY VENOMOUS

Eastern Tiger Snake (*Telescopus semiannulatus*)

Maximum Length: 1 m



Other names: Oostelike Tierslang

A slow-moving nocturnal snake that hunts for lizards, especially geckos, nestling birds, bats and small rodents. It is an excellent climber and spends its days hiding under the bark of trees or under rocks. It can often be seen crossing roads at night. Like the Herald Snake, this snake puts up quite a show when cornered, drawing its head back with coiled neck and striking dramatically with an open mouth. The strikes are rather slow, and the snake may lose its balance. It also flattens the head in the process but does not make a hood. The venom of the Tiger snake is mild and has virtually no effect on humans.







© Johan Marais African Snakebite Institute 2023



Eastern Tiger Snake (Telescopus semiannulatus)

MILDLY VENOMOUS

Average length: 40 - 60 cm. Maximum length: 1 m

Often confused with: Cape Coral Snakes, Harlequin Snakes










MILDLY VENOMOUS

Common Purple-glossed Snake

(Amblyodipsas polylepis)

Maximum Length: 1.12 m



Other names:

Gewone Persglansslang

A common burrowing species that is seldom encountered, except after heavy rains or when soil is excavated. It may also be seen active on overcast days. This snake averages around 60 cm but may exceed a meter in length. If threatened it may hide its head under the body and move the tail tip to distract the aggressor and protect its head.

This snake is easily confused with several other small shiny black snakes, such as the venomous stiletto snakes. It is docile, seldom attempting to bite. The venom of this snake has not been well-studied but is not considered dangerous to humans.







O Johan Marais African Snakebite Institute 2023




Common Purple-glossed Snake (Amblyodipsas polylepis)

MILDLY VENOMOUS

Average length: 40 - 70 cm. Maximum length: 1.12 m










MILDLY VENOMOUS

Olive Grass Snake (*Psammophis mossambicus*)

Maximum Length: 1.8 m



Other names:

Olive Whip/sand Snake, Olyf Grasslang

A large, fast-moving diurnal snake that reaches a maximum length of 1.8 m. It is very quick to escape, invariably disappearing down the closest hole but bites readily if handled. Because of its size and habit of lifting its head well off the ground, this snake is often mistaken for a Black Mamba or female Boomslang. It climbs well and is often found in shrubs. Over much of its range, especially northern KZN, it is one of the most frequently encountered snakes. It is often killed on roads. It preys on lizards, frogs, small mammals, birds and other snakes, including venomous ones. The venom of the Olive Grass Snake is mild and has little effect on humans.







© Johan Marais African Snakebite Institute 2023



Olive Grass Snake (Psammophis mossambicus)


MILDLY VENOMOUS

Average length: 80 cm. Maximum length: 1.8 m

Often confused with: Short-snouted Grass Snakes, Black Mambas, Female Boomslang

Distribution is important

Pale belly








MILDLY VENOMOUS

Western Yellow-bellied Sand Snake

(Psammophis subtaeniatus)

Maximum Length: 1.4 m



Other names:

Westelike Geelpenssandslang

A fast-moving, slender snake that averages 50-80 cm. This snake is active during the day and is without doubt one of our fastest snakes. It is fond of basking but will move off rapidly if disturbed. It is comfortable in trees and shrubs. Like some of the other snakes in the genus *Psammophis*, it will spin if caught by the tail and snap a piece off. The Western Yellow-bellied Sand snake feeds on lizards, frogs, rodents, small birds and other snakes. The venom of this snake is very mild and has no real effect on humans. Sand snakes, Grass snakes and Skaapstekers can be difficult to tell apart – always check the distribution when attempting to identify one of these snakes.







Johan Marais African Snakebite Institute 2023




Western Yellow-bellied Sand Snake (Psammophis subtaeniatus)

MILDLY VENOMOUS

Average length: 80 cm - 1 m. Maximum length: 1.4 m

Often confused with: Short-snouted Grass Snakes, Striped Skaapsteker








CAN INFLICT A NASTY BITE

Southern African Python (*Python natalensis*)

Maximum Length: 6 m



Other names:

Luislang, previously called a Rock Python

The Southern African Python is by far our largest snake, reaching a maximum length of around 6 m and weighing in excess of 65 kg. It feeds largely on warm-blooded prey like small antelope, monkeys, game birds and dassies, but also takes leguaans and even crocodiles. Attacks on humans are rare and fatalities virtually unheard of. This snake is largely active at night but is fond of basking during the day. It is at home in water and can remain submerged for long periods. Females produce 30 – 60 (but in exceptional cases more than 100) eggs, which are roughly the size of a tennis ball. The female remains with her eggs throughout incubation and the young measure 50 – 70 cm in length.







© Johan Marais African Snakebite Institute 2022




Southern African Python (Python natalensis)

NON-VENOMOUS, BUT CAN INFLICT A NASTY BITE

Average length: 3 m. Maximum length: 5.5 m

Often confused with: Puff Adders





SNAKEBITE PROFILE SOUTHERN AFRICAN PYTHON

Python natalensis



Risk of Bites

Pythons are ambush hunters and bites usually occur when walking down pathways or through thick bush, where a python is coiled up awaiting a passing prey animal.



(A) Most Bites


Most bites occur to the lower leg, when the snake lunges at a passerby. They are usually quick to release the bitten person, and often leave a row of small teeth punctures in the leg or calf.



Symptoms & Venom

Pythons have no venom, but have over 80 long recurved teeth, which they use to hang onto prey items. Large pythons may leave nasty wounds after a bite.

See pages 103-105 Snakes and Snakebite in Southern Africa (2024).



First Aid

Immobilise and reassure the patient. Bandage the bitten area to reduce blood loss and transport the victim to a hospital or doctor.



Medical Treatment

In severe bites victims may require stitches. Bites from pythons should be cleaned and disinfected and monitored for signs of infection. Doctors will also administer tetanus injection.








Brown House Snake

(Boaedon capensis)

Maximum Length: 1.5 m



Other names: Bruin Huisslang

The Brown House Snake, true to its name, frequents human dwellings where it feeds largely on rodents and lizards. It is completely harmless and relies on muscle power to constrict its prey. It is widespread in South Africa and very common in suburban gardens.

This snake is active at night and is relatively slow-moving. Most individuals are medium to dark brown with a mother-of-pearl belly and two light lines on either side of the head – one from the nose through the top of the eye to the back of the head and the other from the back of the head and the other from the back of the eye to the angle of the jaw. The Brown House Snake comes in a variety of shades of brown, often with mottling of white along the neck.







© Johan Marais African Snakebite Institute 2022



Brown House Snake (Boaedon capensis)

HARMLESS

Average length: 55 cm. Maximum length: 1.5 m

Can be confused with: Slug-eaters









Common Wolf Snake (Lycophidion capense)

Maximum Length: 65 cm



Other names:

Cape Wolf Snake, Gewone Wolfslang

This common little snake averages 25-40 cm in length and is frequently found in suburban gardens. It is active at night when it hunts for lizards, especially skinks and geckos. The Common Wolf Snake is fond of damp localities where it seeks refuge under rocks, logs and in rubbish heaps. Though harmless it has long, recurved teeth in both the upper and lower jaw, hence the common name. Females produce 3–9 eggs in early summer. The Common Wolf Snake is difficult to distinguish from the venomous stiletto snake. It differs by having a long thin tail, the head is distinct from the body, and it usually has light spots on the back of each body scale.







O Johan Marais African Snakebite Institute 2023




Common Wolf Snake (Lycophidion capense)

HARMLESS

Average length: 40 cm. Maximum length: 64 cm

Often confused with: stiletto snakes










Spotted Bush Snake (*Philothamnus semivariegatus*)

Maximum Length: 1.3 m



Other names:

Variegated Bush Snake; Gespikkelde Bosslang

An alert, day-active snake, that climbs well and is often found between the walls and ceilings of outbuildings, where it hunts for geckos. Sadly, this harmless snake is often mistaken for a Green Mamba or Boomslang and needlessly killed.

The body is bright green to a copper-green with distinct blackish spots on the front half of the snake, but little or no dark marks on the back half. The belly is yellowish to white and the pupil orange.

This snake has keeled belly scales which enables it to climb up the bark of a tree or even up face-brick walls. It is commonly found in suburban gardens and juveniles are often killed by domestic cats.







© Johan Marais African Snakebite Institute 2023



Spotted Bush Snake (Philothamnus semivariegatus)

HARMLESS

Average length: 70 cm. Maximum length: 1.3 m

Often confused with: Western Natal Green Snakes, Green Mambas, Boomslang

Prominent yellow to red iris

Excellent climbers, often found in trees and bushes and in ceilings of outbuildings

In the north of their range, they often have a copper coloured tail

> Yellowish to white belly with keeled ventral and subcaudal scales for climbing

Have bright metallic blue flecks

Double temporal scales



Three upper labials making contact with eye




Blackish spots on the first half of the body





Green Water Snake (Philothamnus hoplogaster)

Maximum Length: 1 m



Other names: South-eastern Green Snake, Groen Waterslang

A smallish bright green snake that averages 25-40 cm in length with a maximum length of 1 m. It is day-active and favours damp localities such as reed swamps, riverine thickets and flood plains where it feeds mainly on frogs. It also eats fish, small lizards and juveniles reportedly eat grasshoppers. Like the harmless natal green snakes, it is often mistaken for a Green Mamba and needlessly killed. Most individuals are bright emerald, green with a whitish belly and often have a yellow snout. Some individuals, especially further north, have a few blackish bars behind the head. This snake is not venomous and completely harmless to humans.







© Johan Marais African Snakebite Institute 2022




Green Water Snake (Philothamnus hoplogaster)

HARMLESS

Average length: 40 - 60 cm. Maximum length: 1 m

Often confused with: Green Mambas, Boomslang, and the Western and Eastern Natal Green Snakes










Common Egg-eater (Dasypeltis scabra)

Maximum Length: 1.16 m



Other names:

Rhombic Egg-eater, Gewone Eiervreter

A common snake throughout most of southern Africa except for true desert and closed canopy forests. It averages around 50 cm and is well adapted at climbing trees. It is most abundant in thornveld and grasslands where it can be found in any situation. It is nocturnal, spending most of the day hiding beneath rocks or under loose bark. When agitated, it will coil and uncoil while striking out viciously. The rubbing scales create a rasping sound similar to a hiss. It is completely harmless, and the teeth are reduced. It feeds exclusively on birds' eggs. It may be confused with the venomous Common Night Adder.







© Johan Marais African Snakebite Institute 2023




Common Egg-eater (Dasypeltis scabra)

HARMLESS

Average length: 55 cm. Maximum length: 1.16 m

Often confused with: Common Night Adders










Zambezi Blind Snake

(Afrotyphlops mucruso)

Maximum Length: 1 m



Other names:

Zambesi se Blindeslang

One of the larger blind snakes that inhabits hard substrates, including termitaria in Mopaneveld and Lowveld regions of northern Limpopo extending into Mozambique and Zimbabwe as well as eastern Botswana. It uses its hard beak to dig into soil and spends most of its life underground, but may surface after rains. Adults are light brown to grey with darker blotches. Females produce more than eight eggs that hatch in just over a month. They feed on termites and their eggs and are preyed upon by a variety of snakes. This snake is harmless and poses no threat to humans. May be confused with Schlegel's Blind Snake but largely the distributions do not overlap.







O Johan Marais African Snakebite Institute 2023

FIRST AID FOR SNAKEBITE



IN THE EVENT OF A SNAKEBITE, DO NOT...

- Do not try to cut or suck out the venom. Never cut the bite site as cutting may do far more harm than good. Sucking does not remove significant amounts of venom.
- Do not apply any electric shock therapy. Shock treatments have been tested extensively and have been shown to have no effect on the outcome of the bite. Electric shocks do not neutralise snake venom.
- ◆ Do not apply a tourniquet. Many people are bitten by harmless snakes or venomous snakes that do not inject sufficient venom to cause serious damage. Arterial or venous tourniquets are not advised in most bites as venom is transported largely through the lymphatic system and not through veins. There is little evidence that a tourniquet could be lifesaving following a snakebite.
- Do not apply heat, ice, lotions or potions. Apart from cleaning the bite site with water and applying a sterile gauze dressing nothing should be applied to the bite. Very hot water does not neutralise snake venom.
- ♣ Do not inject antivenom as a first aid measure. Most snakebite victims do not require antivenom and the polyvalent antivenom is only effective for the bites of certain potentially deadly snakes. If required, antivenom must be injected intravenously by a medical doctor and usually in large quantities. As victims may go into anaphylactic shock when injected with antivenom, adrenaline and other drugs may also be required. Anaphylactic shock is a serious medical condition which may be fatal if not treated promptly.

HRST AD SMILE OF THE SMILE OF T

FIRST AID KITS

We stock a wide range of First Aid for Snakebite Kits, as well as Smart Pressure Bandages, Resuscitators, and first aid books. View the range online here.





FIRST AID FOR SNAKEBITE



IN THE EVENT OF A SNAKEBITE, DO...

- → Do get the victim to a hospital as soon as possible and in a safe manner. This is by far the most important first aid measure. For medical advice in a snakebite emergency, call the African Snakebite Institute +27 (0)82 494 2039.
- → Do keep the victim calm. Often easier said than done as people tend to panic in a snakebite emergency. Many snakebite victims go into shock, experience a drop in blood pressure and may even become nauseous and faint. If possible, do not allow the victim to walk, run, jump etc. Bear in mind that more than 99% of snakebite victims that are hospitalised within a few hours survive their snakebites.
- Do remove jewellery as well as tight clothing. If bitten on the hand/arm/foot/ lower leg, remove rings, bangles, bracelets, watches, anklets and any other tight jewellery including tight clothing.
- → Do administer artificial respiration if patient stops breathing. In serious snakebite cases involving snakes with predominantly neurotoxic venom, like the Black Mamba or Cape Cobra, the victim may soon experience difficulty with breathing, even within 20 minutes. This is a very serious medical condition and assisted breathing could be life-saving. Mouth-to-mouth resuscitation may be beneficial in the short term, but the proper use of a Bag Valve Mask is far better in such an instance.

SPITTING SNAKES

- ★ The most common spitting snake in Southern Africa is the Mozambique Spitting. In the western parts of the region this is replaced by the Black Spitting Cobra and the Zebra Cobra. These snakes spray their venom up to a distance of three metres and do so in self-defence to temporarily blind their attacker so that they can make a quick escape.
- ★ Although spitting snakes accurately aim for the eyes, the venom diffuses into a spray over a wide area and may get onto your hair, face, arms and chest. Venom in the eyes is very painful and must be flushed or diluted as quickly as possible. The production of tears starts this process, but as a first aid measure, flush the eyes with water or a weak (normal) saline solution. The idea is to flush away any excess venom. Transport the victim to a medical doctor, who will examine the eyes for corneal damage and prescribe antibiotic eye drops to prevent secondary infection. There is no reason to use diluted antivenom it has no proven benefits and may cause an allergic reaction.
- If treated quickly, the chance of permanent damage to the eyes is extremely remote.

SNAKEBITE EMERGENCY NUMBERS

Emergency, 0800 3222 911

E-med Rescue 24: 924/ 061 411 600

Crisis Response: 083 3912

African Snakebite Institute: +27 82 494 2039

HOSPITALS AROUND ZIMBABWE

Mater Dei Hospital Tel: +263 29 2240000 -20.180971, 28.596831

Mater Dei Hospital Tel: +263 29 2240000 -20.180971, 28.596831

Victoria Falls Hospital Tel: +263 83 2844262 -17.943546, 25.825903

Kadoma General Hospital Tel: +263 68 2122066 -18.330918, 29.911937 Karanda Hospital Tel: +263 77 408 2211 -16.654871, 31.827439

Mater Dei Hospital Tel: +263 29 2240000 -20.180971, 28.596831

Harare Central Hospital Tel: +263 24 2621100 -17.860487, 311.012865

St. Annes Hospital Tel: +263 24 2303360 -17.788311, 31.034791

SNAKE REMOVERS CONTACTS

Find SNAKE REMOVERS in Zimbabwe

HERE

FREE
ASI SNAKES APP
HERE







The ASI team travels thousands of kilometres a month, often to remote parts of southern Africa, for both snake awareness and venomous snake handling training and for reptile surveys.

We're honoured to have great sponsors, who've made significant changes to our fleet of hardworking vehicles, ensuring they meet our every need.

We'd like to thank our sponsors for making the smooth travel possible.

ASI VEHICLE SPONSORS

























MORE RESOURCES








SHOP OUR RANGE OF BOOKS ONLINE HERE.







CONTACT US

Snake Courses: +27 64 704 7229 or courses@asiorg.co.za

Snake Handling Equipment: +27 60 957 2713 or info@asiorg.co.za

Snake Identification: +27 72 979 1249 or support@asiorg.co.za

Snakebite Emergencies: +27 82 494 2039