COMMON SNAKES OF THE FAST RAND

The East Rand has a great variety of snakes with most of them not considered dangerous. Three species can inflict painful bites, while one is considered deadly.







Rinkhals (Hemachatus haemachatus)



Common Slug-eater (Duberria lutrix lutrix)



Bibron's Stiletto Snake (Atractaspis bibronii) Photo Warren Dick

Black-headed Centipede-eater

(Aparallactus capensis)



Bibron's Stiletto Snake (Atractaspis bibronii)



Bibron's Blind Snake (Afrotyphlops bibronii)



Rhombic Night Adder (Causus rhombeatus)



Rhombic Egg-eater (*Dasypeltis scabra*)



Short-snouted Grass Snake (Psammophis brevirostris)



Brown House Snake




MILDLY

ENOMOUS

Spotted Skaapsteker (Psammophylax rhombeatus)



Aurora House Snake



Cape Wolf Snake (Lycophidion capense capense)



Olive Snake



Herald or Red-lipped Snake (Crotaphopeltis hotamboeia)



Mole Snake

(Boaedon capensis)

(Lamprophis aurora)

(Lycodonomorphus inornatus)


(Pseudaspis cana)

© Jo

JOHAN MARAIS is the author of various books on reptiles including the best-seller *A Complete Guide to Snakes of Southern Africa*. He is a popular public speaker and offers a variety of courses including **Snake Awareness**, **Scorpion Awareness** and **Venomous Snake Handling**. Johan is accredited by the International Society of Zoological Sciences (ISZS) and is a Field Guides Association of Southern Africa (FGASA) and Travel Doctor-approved service provider. His courses are also accredited by the Health Professions Council of South Africa (HPCSA).

Johan Marais | African Snakebite Institute +27 82 494 2039 | johan@asiorg.co.za

www.AFRICANSNAKEBITEINSTITUTE.com



EMERGENCY PROTOCOL

IN THE EVENT OF A SNAKE BITE

- 1 Keep the victim calm, immobilized and transport the victim to the closest hospital without delay.
- 2 If the victim stops breathing, resort to artificial respiration or make use of a Bag Valve Mask.
- **3** Call the Poison Information Centre helpline for further advice: 0861 555 777.

DO NOT

- ... apply a tourniquet.
- ... cut and suck the wound.
- ... use ice or very hot water.
- ... give the victim alcohol.
- ... apply electric shock.
- ... inject antivenom randomly.

Antivenom (*if required*) must be administered by a doctor in a hospital environment.