DANGEROUS SNAKES OF ANGOLA



Angola has over 120 different types of snakes. The majority are not venomous or only mildly venomous. Sixteen can Inflict rather painful bites.

Fifteen species are considered potentially deadly

DANGEROUSHas caused

human fatalities

DANGEROUS


Painful bite, but does not require antivenom




Black Mamba (Dendroaspis polylepis)



Anchieta's Cobra (Naja anchietae)



Brown Forest Cobra (Naja subfulva)



Black-necked Spitting Cobra (Naja nigricollis) - photo Luke Verburgt



Boomslang - male (Dispholidus typus)



Southern Vine Snake
(Thelotornis capensis)



Kunene Coral Snake (Aspidelaps I. cowlesi) - photo Randy Babb






Gaboon Adder (Bitis gabonica)



Puff Adder (Bitis arietans)



Horned Adder (Bitis caudalis)



DANGEROUS


Southern African Python (*Python natalensis*)



Stiletto Snake (Atractaspis sp.) - photo Luke Kemp



Velvety-green Night Adder (Causus resimus)





(Causus bilineatus) - photo Werner Conradie

JOHAN MARAIS is the author of various books on reptiles including the best-seller A Complete Guide to Snakes of Southern Africa. He is a popular public speaker and offers a variety of courses including Snake Awareness, Scorpion Awareness and Venomous Snake Handling. Johan is accredited by the International Society of Zoological Sciences (ISZS) and is a Field Guides Association of Southern Africa (FGASA) and Travel Doctor-approved service provider. His courses are also accredited by the Health Professions Council of South Africa (HPCSA).

Johan Marais | African Snakebite Institute

+27 82 494 2039 | johan@asiorg.co.za





EMERGENCY PROTOCOL

IN THE EVENT OF A SNAKE BITE

1 Keep the victim calm, immobilized and transport the victim to the closest hospital without delay.

2 If the victim stops breathing, resort to artificial respiration or make use of a Bag Valve Mask.

3 Call the Poison Information Centre helpline for further advice: 0861 555 777.

DO NOT —

- ... apply a tourniquet.
- ... cut and suck the wound.
- ... use ice or very hot water.
- ... give the victim alcohol.
- ... apply electric shock.
- ... inject antivenom randomly. Antivenom (if required) must be administered by a doctor in a hospital environment.